

04-2800

A

jc662 U.S. PTO  
04/27/00

## Certificate of Mailing

Date of Deposit April 27, 2000Label Number: EL488650358

I hereby certify under 37 CFR 1.10 that this correspondence is being deposited with the United States Postal Service as "Express Mail Post Office to Addressee" with sufficient postage on the date indicated above and is addressed to: BOX PATENT APPLICATION, Assistant Commissioner for Patents, Washington, D.C. 20231.

Luis A. Cruz

Printed name of person mailing correspondence

Signature of person mailing correspondence



jc530 U.S. PTO  
09/559622

04/27/00

## UTILITY PATENT APPLICATION TRANSMITTAL UNDER 37 CFR §1.53(b)

Attorney Docket Number	01997/521002
Applicant	Rajesh Ranganathan <i>et al.</i>
Title	A NOVEL SEROTONIN-GATED ANION CHANNEL
<b>PRIORITY INFORMATION:</b>	
This application claims priority from U.S. Provisional Application Serial No. 60/131,149, filed April 27, <del>1999</del>	
<b>APPLICATION ELEMENTS:</b>	
Cover sheet	1 page
Specification	47 pages
Claims	4 pages
Abstract	1 page
Drawings (Informals)	22 sheets
Combined Declaration and POA, which is: <input checked="" type="checkbox"/> Unsigned; <input type="checkbox"/> Newly signed for this application; <input type="checkbox"/> A copy from prior application _____ and the entire disclosure of the prior application is considered as being part of the disclosure of this new application and is hereby incorporated by reference therein.	3 pages
Statement Deleting Inventors	0 pages
Sequence Statement	0 pages
Sequence Listing on Paper	0 pages
Sequence Listing on Diskette	0 disks
Small Entity Statement, which is: <input type="checkbox"/> Unsigned; <input type="checkbox"/> Newly signed for this application; <input type="checkbox"/> A copy from prior application _____ and such small entity status is still proper and desired.	0 page

0959522 042700

Preliminary Amendment	0 pages
IDS	0 pages
Form PTO 1449	0 pages
Cited References	0 references
Recordation Form Cover Sheet and Assignment	0 pages
Assignee's Statement	0 pages
English Translation	0 pages
Certified Copy of Priority Document	0 pages
Return Receipt Postcard	1
<b>FILING FEES:</b>	
Basic Filing Fee: \$690.00	\$690.00
Excess Claims Fee: 21 - 20 x \$18.00	\$18.00
Excess Independent Claims Fee: 16 - 3 x \$78.00	\$1,014.00
Multiple Dependent Claims Fee: \$260.00	\$0.00
Total Fees:	\$1,722.00
<input checked="" type="checkbox"/> Enclosed is a check for \$1,722.00 to cover the total fees. <input type="checkbox"/> Charge <b>[**AMOUNT**]</b> to Deposit Account No. 03-2095 to cover the total fees. <input type="checkbox"/> The filing fee is not being paid at this time. <input checked="" type="checkbox"/> Please apply any other charges, or any credits, to Deposit Account No. 03-2095.	
<b>CORRESPONDENCE ADDRESS:</b>	
Kristina Bieker-Brady, Ph.D. Reg. No. 39,109 Clark & Elbing LLP 176 Federal Street Boston, MA 02110	
Telephone: 617-428-0200 Facsimile: 617-428-7045	
 Signature	 Date

Certificate of Mailing


Date of Deposit April 27, 2000

Label Number: EL488650358US

I hereby certify under 37 CFR 1.10 that this correspondence is being deposited with the United States Postal Service as "Express Mail Post Office to Addressee" with sufficient postage on the date indicated above and is addressed to BOX PATENT APPLICATION, Assistant Commissioner for Patents, Washington, D.C. 20231.

Luis A. Cruz

Printed name of person mailing correspondence



Signature of person mailing correspondence

APPLICATION  
FOR  
UNITED STATES LETTERS PATENT

APPLICANT : Rajesh Ranganathan, H.R. Horvitz, and Stephen C. Cannon  
TITLE : A NOVEL SEROTONIN-GATED ANION CHANNEL

A NOVEL SEROTONIN-GATED ANION CHANNEL

Cross Reference to Related Applications

5           This application claims priority from U.S. Provisional Application Serial  
Number 60/131,149, filed April 27, 1999.

Statement as to Federally Sponsored Research

          This research has been funded by NIH Grant R37GM24663. The U.S.  
government has certain rights to the invention.

10                           Background of the Invention

          The invention relates to the diagnosis and treatment of conditions  
associated with serotonin-mediated cellular responses.

          The biogenic amine serotonin plays a role in the modulation of neuronal  
synaptic events as well as non-neuronal cellular signaling. Serotonin acts by  
15   binding to receptors on a variety of cells. These receptors fall into two broad  
functional and structural categories, those acting through G-proteins to mediate  
intracellular signaling, and those that form ion channels. It is generally believed  
that serotonin may act by binding to either G-protein-coupled seven-pass  
transmembrane receptors, or serotonin-gated cation channels. There are six major  
20   classes of G-protein-coupled receptors, each with numerous subtypes. Thus far,  
there is only one class of serotonin-gated ion channels, the 5-HT<sub>3</sub> receptor.  
G- protein-coupled responses can be either excitatory or inhibitory upon activation

by serotonin. Activation of G-protein-coupled receptors by serotonin generally mediates responses which are slower-acting and longer-lasting, while ion channels mediate fast-acting and transitory responses. The 5-HT<sub>3</sub> receptor, comprised of the conducting subunit, 5-HT<sub>3a</sub> and a regulatory subunit, 5-HT<sub>3b</sub> appears to exclusively  
5 elicit excitatory responses that are generally fast-acting and transitory.

The 5-HT<sub>3</sub> receptor is selectively permeable to cations only, such as Na<sup>+</sup> or K<sup>+</sup>, and is very slightly permeable to Ca<sup>2+</sup>. The influx of cations, such as Na<sup>+</sup> into a cell results in depolarization and excitatory neurotransmission. Efflux of cations, such as K<sup>+</sup> hyperpolarizes the cell, thereby reducing the likelihood of  
10 excitation, and generally leads to inhibitory neurotransmission. Since the resting membrane of a typical cell/neuron is much less permeable to Na<sup>+</sup> influx than to K<sup>+</sup> efflux, the opening of a non-selective Na<sup>+</sup>/K<sup>+</sup> channel, such as the 5-HT<sub>3</sub> receptor leads to a dramatic influx of Na<sup>+</sup>, leading to depolarization and excitation.

Serotonin has been implicated in the etiology of many disease states,  
15 including depression, panic disorders, obsessive compulsive disorder, cardiac abnormalities, sleep disorders, eating disorders, nausea and vomiting, gastrointestinal cramps, and migraines. G-protein-coupled serotonin receptors have been implicated in the control of mood (5-HT<sub>1A</sub>), migraine (5-HT<sub>1B</sub>), pain perception (5-HT<sub>1D</sub>), smooth muscle contraction (5-HT<sub>2A</sub>, 5-HT<sub>7</sub>), anxiety (5-HT<sub>2C</sub>), and nausea (5-HT<sub>4</sub>). Activation of the 5-HT<sub>3</sub> receptor by serotonin can  
20 either stimulate or inhibit cardiac function, induce vasodilation, affect lung and intestinal function, cause pain and sensitization of nociceptive neurons, and induce nausea and vomiting. Not surprisingly, many treatments for these disorders are thought to act through serotonergic pathways.

Several classes of drugs thought to modulate the serotonergic pathway exist. For example, selective serotonin re-uptake inhibitors (SSRIs) are used to treat depression. These antidepressants, including Prozac, Zoloft, and Paxil, are believed to act by potentiating serotonin levels at the synapse. Drugs, such as

5    Imitrex, used to treat migraine headaches, act as selective serotonin receptor agonists. Other groups of drugs used to affect mood include monoamine oxidase inhibitors, and selective serotonin receptor antagonists.

While these drugs are administered to humans to treat the above-described disease states, the patients often unpredictably experience a number of

10    side-effects including insomnia, anxiety, chest pain, hypertension, nausea, anorexia, sweating, chills, vomiting, diarrhea, constipation, decreased libido, and abnormal ejaculation. It has been hypothesized that the side-effects result from multiple receptor activation or inactivation when a serotonin agonist or antagonist is given as a treatment. Some of these improperly activated or inactivated

15    receptors may lead to fast- or slow- acting excitatory responses, or slow-acting inhibitory responses, when really only one specific type of response is desired.

A better understanding of serotonin-associated cellular communication could greatly facilitate the discovery of drugs and therapeutic methodologies to treat a broad range of conditions with fewer of the serious and variable side-effects

20    prevalent with currently available drugs that interface with the serotonin pathway. Exactly how the currently available drugs that interface with the serotonin pathway work is not well understood. Agonists, antagonists, and especially serotonin re-uptake inhibitors could affect numerous serotonin receptor subtypes, and the final outcome may be a combined readout of all these varied, and sometimes

25    antagonistic, pathways. It has been hypothesized that the various undesirable side-

effects of a given drug's action is the result of such unwanted activation of the serotonin pathways that are not specific to the condition being treated. Therefore, compounds with a greater specificity for a specific serotonin receptor, for a limited subset of serotonin receptors, or for a specific subtype of a particular class of serotonin receptors would be invaluable to the field of therapeutics for serotonin-mediated disease states.

### Summary of the Invention

We have discovered a serotonin-gated ion channel, MOD-1, that is exclusively permeable to chloride ions, and is not permeable to sodium, potassium, or calcium ions. Activation of this anion channel is most likely to result in an inhibitory response. In some circumstances that are dependent on the reversal potential for chloride (which is a function of the concentration of chloride inside and outside the cell) activation of anion channels could result in an excitatory response. Therefore, it is conceivable that activation of a serotonin-gated anion channel could also result in excitatory neurotransmission.

With the discovery of MOD-1 and the serotonin-gated anion channel that it forms, comes the realization that serotonin may mediate fast-acting, and transitory, inhibitory responses in addition to excitatory responses. It is possible that the activation/inactivation of a MOD-1-like serotonin-gated anion channel, in humans, is associated with some of the effects and/or side-effects of existing serotonin-based drugs. It is also conceivable that many of the serotonin-related diseases are exclusively due to defects in, or associated with, a serotonin-gated anion channel. None of the currently available drugs have been designed to effectively and specifically target this receptor. Therefore, the discoveries of a

serotonin-gated anion channel and the gene that encodes it are invaluable tools for use in discovering diagnostic and therapeutic compounds for the detection and treatment of conditions associated with serotonin-mediated cellular responses.

One way in which a serotonin-gated anion channel can be used as a tool  
5 in drug discovery is by screening existing drugs or drug candidates for their effects on serotonin-gated anion channel activity. Such an experiment can be done using MOD-1 or other serotonin-gated anion channels from non-mammals, such as nematodes, or from lower mammals or humans. Understanding how drugs affect, or do not affect, this anion channel will lend better insight into the overall effect of  
10 a drug's mechanism of action. Also, a better understanding of how this serotonin-gated anion channel is regulated will contribute to a better understanding of how current therapies work. Furthermore, information gained from this screen will permit the development of drugs with higher specificities for a particular type of serotonin-binding receptor that will mediate only the desired response. Such drugs  
15 include those which do not activate a serotonin-gated anion channel, but do activate other serotonin receptors; those that act specifically on a serotonin-gated anion channel, but not on other serotonin receptors; or those that activate a subset of the various serotonin receptors.

Methods of drug discovery are not limited to screening available drugs  
20 only, but also to all compounds and their derivatives that were extracted or synthesized during the development of a given drug affecting serotonin-mediated cellular processes. Furthermore, a *de novo* screen of chemicals can be conducted, with no bias regarding possible functional relevance, for effects on this class of serotonin-gated anion channels.



In a first aspect, the invention features a substantially pure polypeptide that is a serotonin-gated anion channel. In one embodiment, the polypeptide is a substantially pure serotonin-gated anion channel that is permeable to chloride ions. In another embodiment, the polypeptide is MOD-1. In a further embodiment, the polypeptide is a subunit that makes up a multi-subunit serotonin-gated anion channel, permeable to chloride ions. Preferably this polypeptide is from *Caenorhabditis elegans* (*C. elegans*). More preferably this polypeptide is mammalian. Most preferably this polypeptide is human.

In still another embodiment of the invention, the serotonin-gated anion channel is activated by a lower concentration of serotonin than that required to activate the 5-HT<sub>3</sub> receptor. For example, the serotonin-gated anion channel may have a higher binding affinity for serotonin than the 5-HT<sub>3</sub> receptor. This higher affinity can be assessed by calculating and comparing the dissociation constants of serotonin binding to the 5-HT<sub>3</sub> receptor and to a serotonin-gated anion channel.

In another aspect, the invention features a substantially pure nucleic acid sequence encoding a serotonin-gated anion channel. In one embodiment, the substantially pure nucleic acid sequence encodes a serotonin-gated anion channel that is permeable to chloride ions. In another embodiment, the nucleic acid sequence is *mod-1*, and encodes the MOD-1 polypeptide. Preferably this nucleic acid sequence is from *C. elegans*. More preferably the nucleic acid sequence is mammalian. Most preferably the nucleic acid sequence is human.

In another aspect, the invention features an antibody that preferably specifically binds to a serotonin-gated anion channel. In one embodiment, the antibody binds to a serotonin-gated anion channel that is permeable to chloride ions. This invention includes polyclonal, as well as monoclonal antibodies to the

serotonin-gated anion channel.

In another aspect, the invention features a *C. elegans* strain having a mutant *mod-1* gene. In one embodiment of the invention, the *C. elegans* strain has a mutant *mod-1* gene that does not function as a chloride channel. In another  
5 embodiment, the strain has a mutant *mod-1* gene that acts in a dominant-negative manner. In another embodiment, the strain has a mutant *mod-1* gene that encodes a polypeptide that is constitutively active.

In another aspect, the invention features a method for identifying a compound that modulates the biological activity of a serotonin-gated anion  
10 channel. The method includes the steps of: (a) administering a test compound to a serotonin-gated anion channel, and (b) assaying a modulation in the biological activity of a serotonin-gated anion channel. Assaying the modulation of biological activity may be done by measuring the current carried through a channel, or by measuring the amount of serotonin binding to a serotonin-gated anion channel.  
15 The assay can also be a bioassay that involves measuring the rate of locomotion in nematodes having a serotonin-gated anion channel. In one embodiment, the serotonin-gated anion channel is from nematodes. In another embodiment, the serotonin-gated anion channel is from a rat, mouse, or human, and is inserted into a *C. elegans* that either has or does not have a wild-type *C. elegans* serotonin-gated  
20 anion channel. In another embodiment, the serotonin-gated anion channel is a chimeric molecule between the serotonin-gated anion channels from various species. In further various embodiments of this aspect, the channel is in a cell, for example, a neuron or a non-neuronal cell. The channel may also be in a lipid bi-layer, a mammal, or a nematode. In yet another embodiment, the serotonin-gated  
25 anion channel comprises sufficient MOD-1 protein to form a serotonin-gated anion

channel.

In the above aspect of the invention, the test compound is administered in the absence or presence of serotonin, and is administered prior to, simultaneously with, or after administration of serotonin. Administration may also  
5 be in the absence or presence of known drugs that interface with the serotonin pathway. The modulation of the biological activity may be either agonistic or antagonistic. The compound may also be a cell lysate, or isolated from a cell lysate, and may be administered prior to, simultaneously with, or after  
10 administration of serotonin, or any known effector of serotonin-mediated cellular processes.

In another aspect, the invention features a method for treating a condition in a patient by administering an agonist or antagonist of a serotonin-gated anion channel to the patient. Conditions that are treated include migraine headaches, loss of appetite, gain of appetite, insomnia, inability to wake up,  
15 memory loss, inability to learn, nausea and vomiting, gastrointestinal cramps, body temperature deregulation, moods, including depression or mania, abnormal sexual or hallucinogenic behavior, abnormal cardiovascular function, abnormal muscle contraction, and abnormal endocrine regulation.

In another aspect, the invention features a diagnostic probe for  
20 measurement of a serotonin-gated anion channel, either wild-type or mutant, for detecting conditions associated with serotonin-mediated cellular responses. Measurement of a serotonin-gated anion channel includes, but is not limited to detection of nucleic acid levels that code for a serotonin-gated anion channel, levels of a polypeptide that can function as a serotonin-gated anion channel, single  
25 strand confirmation polymorphism analyses, or the flow of current across a

membrane. In one embodiment, the probe is a nucleic acid sequence that encodes a serotonin-gated anion channel, for example, MOD-1, or a polypeptide that is a serotonin-gated anion channel, or an antibody that binds to a serotonin-gated anion channel. In yet another embodiment, the probe is standard electrophysiology voltage clamping equipment that measure the activity of a serotonin-gated anion channel. In still another embodiment, the diagnostic probe is used for pharmacogenetics, i.e., in the analyses of conditions associated with serotonin-mediated cellular responses within an individual, family, or families.

In another aspect, the invention features a method for characterizing drugs associated with serotonin-mediated cellular responses, by measuring serotonin-gated anion channel activity upon drug exposure. The drugs include those already currently available for the treatment of serotonin-mediated responses, as well as small molecules that are similar in structure to these drugs. This invention also includes the discovery of any compounds which have not yet been identified as therapeutic for serotonin-mediated cellular responses.

In another aspect, the invention features a method for decreasing serotonin-gated anion channel function by decreasing the level of a serotonin-gated anion channel polypeptide with antisense RNA to the serotonin-gated anion channel, and the antisense RNA itself. Preferably the level of the serotonin-gated anion channel is decreased at least 25%, more preferably at least 50%, 70%, or 80%, and most preferably at least 95%, compared to a control (e.g., a serotonin-gated anion channel that is not contacted with an antisense RNA, or that is contacted with a nonsense RNA sequence). In addition, preferably the antisense RNA is antisense *mod-1* RNA. Such nucleic acids of the invention and methods for using them may be identified according to a method involving: (a) providing a

cell sample; (b) introducing by transformation into the cell sample, a test nucleic acid sequence for a serotonin-gated anion channel; (c) expressing the test nucleic acid of a serotonin-gated anion channel within the cell sample; and (d) determining whether the cell sample exhibits altered serotonin-gated anion channel activity, whereby either increased or decreased channel activity identifies a nucleic acid which may be used to alter serotonin-gated anion channel function. Preferably the cell is a non-neuronal cell. Most preferably the cell is a neuronal cell.

In another aspect, the invention features a method for decreasing the function of a serotonin-gated anion channel by administering an antibody that specifically binds to a serotonin-gated anion channel, or binds to a peptide from that channel. This method includes, but is not limited to, using an antibody to MOD-1 as the antibody, and a channel formed by any MOD-1 polypeptide as the channel whose function is inhibited. The methods also includes using a mammalian antibody to decrease the function of a mammalian serotonin-gated anion channel. This method also includes administration of the antibody *in vivo* or *in vitro*.

In another aspect, the invention features a method for modulating serotonin-gated anion channel activity using a nucleic acid vector encoding a serotonin-gated anion channel, and administering enough vector to alter activity of the serotonin-gated anion channels of at least one cell. In one embodiment, the vector is operably linked to a promoter. In other embodiments, the vector encodes a wild-type serotonin-gated anion channel or a mutant serotonin-gated anion channel. The mutant serotonin-gated anion channel may include, but is not limited to, a mutant channel that is a loss-of-function mutant, a dominant negative mutant, or a constitutively active mutant. In a preferred embodiment, the nucleic acid

vector encodes a wild-type or mutant MOD-1 polypeptide. In other embodiments, administration may occur *in vitro* or *in vivo*. In further embodiments, the vector encodes a polypeptide that affects the function of a serotonin-gated anion channel, or that affects molecules that are targeted subsequent to activation of a serotonin-gated anion channel. These molecules include, but are not limited to, protein kinases, protein phosphatases, and proteases.

In another aspect, the invention features a method for testing a patient having a serotonin-mediated condition, for his/her pre-disposition to respond to therapy, or to experience side-effects due to administration of a specific therapy.

10 The method comprises:

- (a) determining the characteristics of a serotonin-gated anion channel from tissues of the patient, where the characteristics are indicative of the abnormal activity of a serotonin-gated anion channel; and
- (b) administering to the patient a suitable therapeutic agent relative to the degree of abnormal activity of the serotonin-gated anion channel in step (a).

In various embodiments, the abnormal activity is due to mutations in a serotonin-gated anion channel protein, altered levels of synthesis of mRNA of a serotonin-gated anion channel, altered serotonin-gated anion channel protein levels in tissues. In another embodiment, the method further comprises characterizing other serotonin-mediated receptors in the patients. Therapeutic agents to be used in accordance with the present invention may be selected from the group consisting of inhibitors or activators of serotonin-mediated pathways, including therapeutics which are currently available, as well as those which are discovered,

as described herein.

In accordance with the present invention, there is provided a method for the identification of a patient possessing a serotonin-mediated condition to be responsive to therapies for the condition, or to experience side-effects due to administration of a specific therapy. This method comprises:

(a) determining the characteristics of a serotonin-gated anion channel gene allele in a biological sample of the patient directly, using appropriate probes to a serotonin-gated anion channel, or indirectly, by phenotyping, and;

(b) correlating the genotype or phenotype with appropriated drug and/or dosage.

The presence or absence of a specific serotonin-gated anion channel allele indicates a predisposition, or lack thereof, to respond to serotonin-mediated therapies.

In another aspect, the invention claims a method for identifying a gene that is structurally related to a serotonin-gated anion channel. This method involves identifying a gene by designing probes or primers, including degenerate oligonucleotides, to specific sequences. These primers or probes encode structurally significant amino acid sequence (e.g., the sequence that forms the transmembrane portions of the serotonin-gated anion channel), and are used to screen large genomic or cDNA libraries. If a PCR approach is utilized, the primers are optionally designed to allow cloning of the amplified product into a suitable vector. PCR is particularly useful for screening cDNA libraries from rare tissue types. The method also includes identifying a gene using antibodies, using nucleic

acid or amino acid scanning databases and computer programs, and screening for genes that function in a manner similar to, or different from, a serotonin-gated anion channel.

In a related aspect, the invention features the nucleic acid sequence  
5 identified by the method of identifying a gene which is structurally related to a serotonin-gated anion channel. This gene may be isolated from nematodes or mammals, preferably from rodents, and most preferably from human.

In another aspect, the invention features a transgenic, or other mutant  
10 animal, that over-expresses or under-expresses a serotonin-gated anion channel, or expresses a dominant-negative serotonin-gated anion channel. The invention includes a nematode or a non-human mammal, for example, a mouse, as the animal. In one embodiment, the animal over-expresses a serotonin-gated anion channel that is constitutively active.

In two additional aspects, the invention features a transgenic animal and  
15 methods of using the animal for the detection of therapeutics for conditions associated with serotonin-mediated cellular responses. Preferably the animal over-expresses a serotonin-mediated anion channel polypeptide, either wild-type or mutant, or expresses an antisense RNA to a serotonin-gated anion channel or a serotonin-gated anion channel fragment. In one embodiment, the animal also has a  
20 genetic predisposition to conditions associated with serotonin-mediated cellular responses.

In yet another aspect, the invention features a method for identifying a compound that modulates the activity of a serotonin-gated anion channel by exposing a nematode to a test compound, assaying the rate of locomotion, and  
25 comparing the locomotion rate to that of a nematode receiving no test compound,



serotonin, or a placebo, where a modulation in the rate of locomotion indicates a compound that modulates the activity of a serotonin-gated anion channel. The test compound may be applied at various concentrations. In addition, the nematodes used in the screen may be bacterial-lawn deprived prior to beginning the screen.

5 In still another aspect, the invention features a method for identifying a compound that modulates the activity of a serotonin-gated anion channel. The method involves exposing a nematode to test compound, quantifying the number of nematodes actively swimming after exposure to the compound, and comparing that number to that of a control receiving no test compound, serotonin, or a  
10 placebo, where a modulation in the number of actively swimming nematodes indicates a compound that modulates the activity of a serotonin-gated anion channel. The test compound may be applied at various concentrations. In addition, the nematodes used in the screen may be bacterial-lawn deprived prior to beginning the screen.

15 By "treatment" is meant the submission or subjection of an animal, cell, lysate or extract derived from a cell, or molecule derived from a cell to a test compound or stimulus to a serotonin-mediated response.

By a "test compound" is meant a chemical, be it naturally-occurring or artificially-derived, that is surveyed for its ability to modulate serotonin-mediated  
20 cellular responses, by employing one of the assay methods described herein. Test compounds may include, for example, peptides, polypeptides, synthetic organic molecules, naturally occurring organic molecules, nucleic acid molecules, and components thereof.

By "treat" is meant to submit or subject an animal, cell, lysate or extract  
25 derived from a cell, or molecule derived from a cell to a test compound or

stimulus to a serotonin-mediated response.

By a "substantially pure polypeptide" is meant a polypeptide that has been separated from the components that naturally accompany it. Typically, the polypeptide is substantially pure when it is at least 60%, by weight, free from the proteins and naturally-occurring organic molecules with which it is naturally associated. Preferably the polypeptide is a serotonin-gated anion channel polypeptide that is at least 75%, more preferably at least 90%, and most preferably at least 99%, by weight, pure. A substantially pure serotonin-gated anion channel polypeptide may be obtained, for example, by extraction from a natural source (e.g., a neuronal or smooth muscle cell), by expression of a recombinant nucleic acid encoding a serotonin-gated anion channel polypeptide, or by chemically synthesizing the protein. Purity can be assayed by any appropriate method, e.g., by column chromatography, polyacrylamide gel electrophoresis, agarose gel electrophoresis, optical density, or HPLC analysis.

A protein is substantially free of naturally associated components when it is separated from those contaminants which accompany it in its natural state. Thus, a protein which is chemically synthesized or produced in a cellular system different from the cell from which it naturally originates will be substantially free from its naturally associated components. Accordingly, substantially pure polypeptides include those derived from eukaryotic organisms but synthesized in *E. coli* or other prokaryotes.

By a "purified nucleic acid" is meant a nucleic acid that is free of the genes which, in the naturally-occurring genome of the organism from which the nucleic acid of the invention is derived, flank the gene. The term therefore includes, for example, a recombinant DNA that is incorporated into a vector, into

an autonomously replicating plasmid or virus, or into the genomic DNA of a prokaryote or eukaryote, or which exists as a separate molecule (e.g., a cDNA or a genomic or cDNA fragment produced by PCR or restriction endonuclease digestion) independent of other sequences. It also includes recombinant DNA that  
5 is part of a hybrid gene encoding additional polypeptide sequence.

By a “serotonin-gated anion channel” is meant a channel whose opening is regulated by serotonin binding to the channel. The opening of the channel selectively permits passage of anions from one side of the channel to the other. In one embodiment, the anion is chloride. Preferably the nucleic acid sequence  
10 encoding a serotonin-gated anion channel hybridizes to a *mod-1* nucleic acid sequence.

By a “*mod-1* gene” is meant a gene encoding a polypeptide that is a serotonin-gated anion channel. In one embodiment, the *mod-1* gene is from *C. elegans*.

By a “MOD-1 protein” or “MOD-1 polypeptide” is meant a polypeptide or fragment thereof, encoded by the *mod-1* gene. In one embodiment, the MOD-1 protein or polypeptide is from *C. elegans*.  
15

By “specifically binds” is meant an antibody that recognizes and binds to a serotonin-gated anion channel, but which does not substantially recognize and  
20 bind to other molecules in a sample, e.g., a biological sample that naturally includes other proteins.

By “mutant” is meant different from what normally appears, occurs, or functions. As used herein, the term refers to a nucleic acid sequence that is different from the wild-type sequence. This term also describes a protein encoded  
25 by the mutant nucleic acid sequence. The term also means an organism that

contains a mutant nucleic acid sequence.

By “biological activity” is meant functional events mediated by a protein. In some embodiments, this includes events assayed by measuring the influx of ions into or out of a cell, or assaying the amount of serotonin binding to a channel. It also includes interactions of a polypeptide with another polypeptide. It also includes events that modify behavior or behavioral states. Such behavior includes, but is not limited to, movement, sexual behavior, or hallucinogenic behavior. Such behavioral states include, but are not limited to, migraine headaches, loss of appetite, gain of appetite, insomnia, inability to wake up, memory loss, nausea or vomiting, gastrointestinal cramps, ability or inability to learn, body temperature deregulation, moods, such as depression or mania, abnormal cardiovascular function, abnormal muscle contraction, and abnormal endocrine regulation.

As used herein, by “modulates” is meant increasing or decreasing the biological activity. Preferably the biological activity is increased or decreased 10% relative to a control. More preferably the biological activity is increased or decreased 50% relative to a control. Most preferably the biological activity is increased or decreased 90% relative to a control.

By “assaying” is meant analyzing the effect of a treatment or exposure, be it chemical or physical, administered to a whole animal or cells derived therefrom. The material being analyzed may be an animal, a cell, a lysate or extract derived from a cell, or a molecule derived from a cell. The analysis may be, for example, for the purpose of detecting current flow across a membrane, the rate of locomotion of an animal, altered gene expression, altered nucleic acid stability (e.g., mRNA stability), altered protein stability, altered protein levels, or

altered protein biological activity. The means for analyzing may include, for example, recording current changes under voltage-clamp, voltage changes in current-clamp, or extracellular potentials, counting movements of an animal under a dissecting microscope, nucleic acid amplification techniques, reporter gene  
5 assays, antibody labeling, immunoprecipitation and phosphorylation assays, and other techniques known in the art for conducting the analyses of the invention.

By "neuron" is meant a cell of ectodermal embryonic origin derived from any part of the nervous system of an animal. Neurons express well-characterized neuron-specific markers that include neurofilament proteins, MAP2,  
10 and class III  $\beta$ -tubulin. Included as neurons are, for example, hippocampal, cortical, midbrain dopaminergic, motor, sensory, sympathetic, septal cholinergic, and cerebellar neurons.

As used herein, by "measuring" is meant assessing an anion channel activity. Measuring can be done by use of standard electrophysiology voltage-  
15 clamping or patch-clamping equipment.

By "condition" is meant a state of being or feeling. Conditions include, but are not limited to, migraine headaches, loss of appetite, gain of appetite, insomnia, inability to wake up, memory loss, nausea or vomiting, gastrointestinal cramps, ability or inability to learn, body temperature deregulation, moods such as  
20 depression or mania, abnormal sexual or hallucinogenic behavior, abnormal cardiovascular function, abnormal muscle contraction, and abnormal endocrine regulation.

By "promoter" is meant a minimal sequence sufficient to direct transcription of an operably-linked gene.

By “operably linked” is meant that a gene and one or more regulatory sequences are connected in such a way as to permit gene expression when the appropriate molecules (e.g., transcriptional activator proteins) are bound to the regulatory sequences.

5 By “dominant-negative” is meant a nucleic acid sequence encoding a polypeptide which when expressed acts in a way to inhibit another polypeptide. This term also refers to the polypeptide itself. In one embodiment, the polypeptide that is inhibited is a wild-type polypeptide, and the dominant negative sequence encodes a mutant polypeptide of the same gene.

10 As referred to herein, by “constitutively active” is meant a nucleic acid sequence that encodes a polypeptide, which when expressed is in an active form at least as, or more often as the wild-type polypeptide is, in a cell in which wild-type polypeptide is naturally expressed. The polypeptide may be in an active form by being phosphorylated, or dephosphorylated, or cleaved from a propeptide to a  
15 peptide, or being ligand independent, or being mutated.

By “transgenic” is meant any cell or organism that includes a DNA sequence (transgene) that is inserted by artifice into a cell and becomes part of the genome of the organism that develops from that cell. As used herein, the transgenic organism is generally a transgenic non-human mammal (e.g., rodents  
20 such as rats or mice) or invertebrate (e.g., *Caenorhabditis elegans*).

By “antisense” is meant a nucleic acid sequence, regardless of length, that is complementary to the coding strand gene encoding a serotonin-gated anion channel. Preferably the antisense nucleic acid is capable of decreasing the activity of a serotonin-gated anion channel when present in a cell that normally is  
25 modulated by serotonin. Preferably the decrease is at least 10%, relative to a

control, more preferably 25%, and most preferably 95%.

By “expose” is meant to allow contact between an animal, cell, lysate or extract derived from a cell, or molecule derived from a cell, and a test compound or activator of a serotonergic response.

5 By “characteristics” is meant properties or features. Characteristics include, but are not limited to, the nucleic acid sequence of a gene, or various alleles of a gene, the amino acid sequence of a protein, the level of expression of proteins or mRNA, and altered protein levels in tissues.

10 By “inhibit” is meant to decrease the level of expression of a serotonin-gated anion channel, or to decrease the function or activity of a serotonin-gated anion channel. Preferably the expression, function, or activity of the channel is decreased at least 25%, more preferably at least 50%, 70%, or 80%, and most preferably at least 95%, compared to a control (e.g., one which is not contacted with a test compound or an antisense nucleic acid).

15 Brief Description of the Drawings

Fig. 1 shows the genomic sequence of *C. elegans mod-1*.

Fig. 2 shows the cDNA sequence encoding the *C. elegans* MOD-1 polypeptide.

Fig. 3 shows the *C. elegans* MOD-1 predicted amino acid sequence.

20 Fig. 4 shows the structure of the *C. elegans mod-1* cDNA.

Fig. 5 is a map of the *C. elegans* strain carrying the ok103 mutation at the *mod-1* locus. The ok103 mutation is a 4135 base pair deletion in the *mod-1* genomic locus.

Fig. 6 shows the genomic sequence of the *C. elegans mod-1* gene with the ok103 mutation.

Fig. 7 shows the genomic sequence of the *C. elegans mod-1* gene with the n3034 mutation.

5 Fig. 8 shows the cDNA sequence of the *C. elegans mod-1* gene with the n3034 mutation.

Fig. 9A is a time trace of the MOD-1 channel activated by serotonin.

Fig. 9B is a graph illustrating a dose response curve of MOD-1 channel activity with respect to serotonin concentration.

10 Fig. 9C is a graph illustrating the peak ionic current conducted by the MOD-1 channel in response to a variety of ligands, including acetylcholine, GABA, glycine, Q-107 (quipazine dimaleate, a 5-HT<sub>3</sub> receptor agonist), and serotonin.

Fig. 10 is a graph illustrating the selectivity the MOD-1 channel has for chloride ions. As the concentration of chloride ions outside of the cell increases, the reversal potential becomes more negative.

15 Fig. 11 is a time trace of a MOD-1-like response from rat brain RNA.

### Detailed Description of the Invention

#### MOD-1 is associated with locomotion in *C. elegans*

20 Hermaphrodites respond to the presence of a bacterial lawn (their food source) by slowing their rate of locomotion. Animals deprived of bacteria for 30 minutes exhibit enhanced slowing when they encounter a bacterial lawn. Genetic, pharmacological, and laser ablation studies demonstrate that this modulatory



response is mediated, in part, by serotonin.

Nematodes to be tested in the locomotion assay were picked as L4 animals 16-20 hours prior to the assay. Locomotion was assayed by placing the nematode on an assay plate (prepared by spreading a solution of *E. coli* strain HB101 on NGM agar in 5 cm plates, using a ring with an inner diameter of approximately 1 cm and an outer diameter of approximately 3.5 cm, and allowing the bacteria to grow 13-15 hours at 37°C), observing each nematode under a dissecting microscope for 20 seconds, and counting the number of dorsal-ventral bends that occur in the anterior portion of the body during the interval.

For satiated animals, locomotion rate was assayed by removing five animals from plates with ample bacteria, washing them in S-basal buffer, and transferring them to the clear zone of the bacterial lawn of an assay plate using a capillary pipette. Beginning five minutes after transfer, the number of body bends was counted, as described above. This procedure was performed for each of the 5 animals.

To assay locomotion rates in food-deprived animals, 5-15 animals were removed from plates with ample food, washed twice in S-basal buffer, and transferred to 5 cm NGM agar plates without bacteria. Food-deprived animals were incubated on these plates for thirty minutes at room temperature, and then were transferred to assay plates. Locomotion was assayed, as described above for satiated animals.

The *mod-1* mutant, obtained from the screen, as described above, exhibited a dominant phenotype of lack of decreased locomotion after being deprived of bacteria and then returned to a bacterial lawn.

MOD-1 is associated with locomotion seen in *C. elegans* exposed to serotonin

Animals to be tested for sensitivity or insensitivity to serotonin in this liquid locomotion assay were picked as L4 animals 16-20 hours prior to assay and the plates were coded so that the experimenter was blind to the genotype of the animals to be scored. On the day of the assay, serotonin (as a creatinine sulphate salt) was dissolved in M9 (at the required concentrations) just before use, and 200  $\mu$ l were aliquoted to the wells of a flat-bottomed 96-well polystyrene plate.

Twenty nematodes of each genotype to be tested were then transferred from the plate into a well containing liquid, using bacteria as glue. Care was taken not to hurt the animals while dislodging them from the pick, and the animals were observed immediately after transfer to ensure that all of them began to exhibit swimming motions. Then at various time intervals, the number of nematodes in each well that were still actively swimming was quantified.

The *mod-1* mutants, as described in the previous section, were further characterized using this technique. Animals carrying the n3034 mutation exhibited a dominant phenotype of insensitivity to exogenous serotonin in liquid locomotion assays. Animals carrying the ok103 mutation exhibited a recessive phenotype of insensitivity to exogenous serotonin in liquid locomotion assays.

The following examples are provided to illustrate the invention. These examples should not be construed as limiting.

Example 1: Cloning of the *mod-1* gene

Both the wild type, and mutant *mod-1* cDNA have been obtained. The dominant serotonin resistance phenotype of animals carrying the *mod-1*(n3034) allele was used to genetically map *mod-1*(n3034) to a 0.7 map-unit interval on

chromosome V. Deficiency analysis showed that the dominant serotonin resistance phenotype is not due to a haploinsufficiency of the *mod-1* locus. The recessive nature of the serotonin resistance phenotype at early time points was exploited to perform standard transformation rescue experiments, and  
5 subsequently, the gene was cloned.

#### Example 2: MOD-1 encodes a ligand-gated ion channel

The protein encoded by the *mod-1* open reading frame responsible for the rescue is structurally similar to ligand-gated ion channels that belong to the nicotinic acetylcholine receptor (nAChR) family. The nAChR family members are  
10 all pentameric channels with large N-terminal extracellular ligand-binding domains, four highly conserved transmembrane domains (M1-M4), and relatively divergent cytoplasmic domains between M3 and M4. nAChR family members include channels gated by acetylcholine, glycine, GABA, avermectin, and serotonin. Within the members of the nAChR family, structure-function analysis  
15 has been performed primarily on the acetylcholine receptor, but many structural and functional parallels have been seen with the other family members as well. In addition, chimeric channel studies show that there is a great deal of conservation at the functional level, even across the different ligand-gated members of the family. The M2 domains of the various subunits are predicted to line the pore of the  
20 channels. Site-directed mutagenesis studies of residues within this domain have demonstrated that ion specificity and modulation of the magnitude and frequency of current flux are determined, at least in part, by the residues that line the pore and those that are immediately adjacent to the pore on both the extracellular and cytoplasmic sides. Based on primary sequence analysis, MOD-1 appears to be

equally divergent from all cloned nAChR family members.

Example 3: *mod-1* forms a serotonin-gated anion channel

MOD-1 was heterologously expressed in *Xenopus* oocytes, injected with 50 nl of *C. elegans* RNA, or MOD-1 was expressed in HEK cells transiently  
5 transfected by calcium phosphate precipitation. Forty-eight to 72 hours later, the cells or oocytes were screened under a voltage clamp. Application of 100 nM serotonin elicited large inward currents at a holding potential of -70 mV. Uninjected oocytes and nontransfected cells had no response to 10  $\mu$ M serotonin. Application of 1 mM of other agonists of ligand-gated ion channels, such as  
10 acetylcholine, GABA, or glycine elicited little or no response from the MOD-1 channel.

Example 4: MOD-1 is permeable to chloride ions

Ion selectivity was determined by measuring changes in the reversal potential (voltage at which the serotonin response changes from an inward,  
15 negative, to an outward, positive, current) in response to varying the ionic composition of the bath solution. The reversal potential was insensitive to changes in cations ( $\text{Na}^+$  or  $\text{K}^+$ ), but shifted by approximately 50 mV for each 10-fold change in extracellular chloride concentration.

Example 5: Antibodies to MOD-1

20 Using MOD-1 polypeptides described herein, anti-MOD-1 antibodies were produced using standard techniques. Peptides to either the putative N-terminal extracellular domain, or to a putative C-terminal intracellular domain,

located between transmembrane-spanning regions III and IV, were synthesized, and coupled to GST using standard techniques. The peptides were used to immunize two different rabbits and three different rats. The antibodies were then affinity purified on a HIS-tagged-MOD-1 affinity column using standard techniques, and were shown to specifically identify GST-MOD-1, by Western blotting.

Polypeptides for additional antibody production may be produced by recombinant or synthetic peptide techniques (see, e.g., Solid Phase Peptide Synthesis, 2nd ed., 1984 The Pierce Chemical Co., Rockford, IL; Ausubel et al., *supra*).

Alternatively, monoclonal antibodies may be prepared using a serotonin-gated anion channel polypeptide (or immunogenic fragment or analog) and standard hybridoma technology (see, e.g., Kohler et al., Nature 256:495, 1975; Kohler et al., Eur. J. Immunol. 6:511, 1976; Kohler et al., Eur. J. Immunol. 6:292, 1976; Hammerling et al., In *Monoclonal Antibodies and T Cell Hybridomas*, Elsevier, NY, 1981; Ausubel et al., *supra*).

Once produced, polyclonal or monoclonal antibodies are tested for specific recognition by Western blot or immunoprecipitation analysis (by the methods described in Ausubel et al., *supra*). Antibodies that specifically recognize a serotonin-gated anion channel polypeptide, as described herein, are considered to be useful in the invention. Anti-serotonin-gated anion channel antibodies, as isolated above, may be used, e.g., in an immunoassay to assay or monitor the level of a serotonin-gated anion channel polypeptide produced by *C. elegans* or a mammal, or to screen for compounds that modulate serotonin-gated anion channel polypeptide production. Anti-serotonin-gated anion channel antibodies may also

be used to identify cells that express a serotonin-gated anion channel gene.

Example 6: Cloning of mammalian serotonin-gated chloride channels with biophysical properties similar to the *C. elegans* serotonin-gated anion channel

Based on our isolation of a novel nematode *mod-1* cDNA, the isolation  
5 of mammalian nucleic acid sequences encoding a serotonin-gated anion channel, including human sequences, is made possible using the strategies described herein and standard techniques.

I. Expression cloning using *Xenopus* oocytes

A. Poly(A)+ RNA isolated from various tissues types is injected into *Xenopus*  
10 oocytes. Initial screening is performed on total Poly(A)+ RNA from human brain, human spinal cord, human testes, rat brain, rat testes, mouse brain, and mouse testes.

B. Functional expression of serotonin-gated ion channels is assayed by measuring  
ion current in voltage-clamped oocytes elicited by application of serotonin at a  
15 concentration of 100 nM or less. Several strategies can be used to identify those serotonin-gated currents that are likely to be conducted by an anion-selective ionotropic channel, comparable to MOD-1.

1) High affinity for serotonin. We have found that MOD-1 channels  
20 are activated by lower concentrations of serotonin than 5-HT<sub>3a</sub> type serotonin receptors. MOD-1 has a half-maximal response with 50 nM serotonin. For 5-HT<sub>3a</sub> channels, no response is detectable in

1  $\mu$ M serotonin, but 10  $\mu$ M serotonin elicits a robust current. The lower affinity of the 5-HT<sub>3a</sub> receptor for serotonin is also reflected in its faster and more complete "washout" (reduction of current back to baseline when the drug is washed off) compared to MOD-1.

5 Serotonin-gated anion channels that have a lower affinity for serotonin can also be isolated using the methods, as described above.

10 2) Shifts in the reversal potential caused by changes in [Cl]<sub>o</sub>. In standard external saline (140 mM NaCl, 2 mM CaCl<sub>2</sub>, 1 mM MgCl<sub>2</sub>, 2.5 mM glucose, and 10 mM HEPES) the reversal potential (E<sub>rev</sub>) for MOD-1 is about -20 mV, whereas the 5-HT<sub>3a</sub> current reverses at about 0 mV. Decreasing the extracellular Cl<sup>-</sup> to 50 mM by substitution with gluconate, while keeping the cation concentration fixed, will cause E<sub>rev</sub> of the MOD-1 current to shift to > +15 mV  
15 whereas E<sub>rev</sub> for the 5-HT<sub>3a</sub> current will not be changed.

20 3) Exclusion of metabotropic (second messenger dependent) serotonin-gated currents. Ligand binding to metabotropic 5-HT receptors, such as the 5-HT<sub>1C</sub> receptor, expressed in *Xenopus* oocytes elicits a large Cl<sup>-</sup> current due to IP<sub>3</sub>-triggered release of intracellular Ca<sup>2+</sup> and subsequent activation of endogenous Ca<sup>2+</sup>-activated Cl<sup>-</sup> channels. This signaling pathway can be suppressed by direct injection of IP<sub>3</sub>, which exhausts the stores of releasable Ca<sup>2+</sup> and thereby causes the oocyte to be refractory to metabotropic 5-HT

receptor responses that depend on internal  $\text{Ca}^{2+}$  release. Serotonin-gated responses acting through second messenger systems also differ from ionotropic responses in that second-messenger based systems often give rise to oscillations in ion current. These oscillations arise from the dynamic aspects of intracellular  $\text{Ca}^{2+}$  regulation. In contrast, current conducted by ionotropic receptors always have a monophasic time course.

- 4) 5-HT<sub>3a</sub> specific toxin. The 41-amino acid snail venom  $\sigma$ -conotoxin GVIIIA is a potent competitive antagonist of the 5-HT<sub>3a</sub> channel (IC<sub>50</sub> on the order of 50 nM). One can test whether MOD-1, like all other known 5-HT class receptors, is insensitive to  $\mu\text{M}$  concentrations of this conotoxin. This strategy may provide a simple method for eliminating the only other known serotonin-gated ionotropic receptors (5-HT<sub>3a</sub> and 5-HT<sub>3b</sub>).
- 5) Suppression of all known forms of serotonin channels. Ideally, expression cloning of mammalian serotonin-gated chloride channels is achieved under conditions in which all other known serotonin gated channels and receptors are blocked. The blockage of these serotonin gated channels and receptors can be accomplished by bathing the oocyte in compounds that inhibit their function. For example, the oocytes may be bathed in a solution containing the calcium chelator BAPTA-AM. The concentration of BAPTA-AM in the bath is approximately 200  $\mu\text{M}$ . While BAPTA-AM blocks other known serotonin gated receptors, it does not



affect the function of MOD-1.

Alternatively, the bath may contain a high concentration of calcium, for example, approximately 10 mM, that is sufficient to block serotonin gated cation channels, but does not block MOD-1. In another method, serotonin gated  $\text{Na}^+/\text{K}^+$  channels may be blocked using, for example, choline chloride or N-methyl glucamine. In other methods, the bath may contain, specific serotonin antagonists. For example, granisetron or ondansetron (at approximately 1  $\mu\text{M}$  concentrations) may be used to inhibit the serotonin-gated cation channel.

One skilled in the art can appreciate that a combination of these methods may provide the maximal blockage of all known serotonin gated ion channels and receptors. For example, one preferred combination of serotonin gated channel and receptor blockers is BAPTA-AM and/or choline chloride or N-methyl-glucamine.

A method to identify a mammalian serotonin-gated chloride channel was carried out essentially as described above. Oocytes were injected with rat cortex RNA. Six days later, the oocytes were bathed in BAPTA-AM (200  $\mu\text{M}$ ) for 2 hours at room temperature, to deplete internal  $\text{Ca}^{++}$  stores, thereby preventing G-protein-coupled serotonin receptors from having an effect through calcium-dependent second messenger pathways. The oocytes were then screened under a voltage clamp, in a buffer containing 96 mM NaCl, 2 mM KCl, 0.3 mM  $\text{CaCl}_2$ , 1 mM  $\text{MgCl}_2$ , and 5 mM HEPES (pH 7.5), and with 100 mM Tris added. Addition of Tris completely blocks the  $5\text{HT}_3$  cation channel. Application of 1  $\mu\text{M}$  serotonin

for 1 minute elicited inward currents at a holding potential of -70 mV in about 10% of the oocytes (Fig. 11).

These results indicate that rat brain RNA elicits a MOD-1-like serotonin-activated current from oocytes. This current is not a 5HT<sub>3a</sub> (and is very unlikely to be another permeant 5-HT ionotropic cation channel receptor) or a G-protein-coupled serotonin receptor-activated current. The isolation of the specific RNA that mediates these results is achieved, as described below.

C. When a response that fits some or all of the criteria outlined above has been identified, one can size fractionate the RNA as the first step toward isolating a specific RNA (and corresponding cDNA). Our preliminary studies of MOD-1 and data from the cloning of the 5-HT<sub>3a</sub> cation channel demonstrate that functional serotonin-gated channels can be formed by homomultimers of a single gene product. This result tremendously improves the feasibility of the expression-cloning strategy. A single RNA species is capable of coding for a functional ionotropic channel, which implies that RNA size fractionation should not cause a loss of serotonin response, as might occur with a heteromultimeric channel protein. Sucrose gradients are used to size fractionate total RNA and individual fractions are injected into oocytes to test for serotonin-gated responses. Once a small enough pool has been determined by such methods, the RNA from that pool is used to prepare cDNA libraries in appropriate vectors. Resulting clones are end-sequenced, and RNA is synthesized from distinct clones and tested in oocytes for the required activity. Clones that produce the desired response are then sequenced in their entirety.

0959632 04300  
002240 2296550

If the results from these experiments indicate that the mammalian serotonin-gated anion channel consists of a heteromultimer rather than a homomultimer, then various pools of mammalian RNA are injected into the oocyte along with any RNAs encoding putative serotonin-gated anion channels. The functional expression of serotonin-gated anion channels can be assayed, as described above. When a pool meets the requirements of contributing to the make-up of a serotonin-gated anion channel, as outlined above, the pool can be further fractionated, and the assays, as described above, are repeated. The methods used to identify and clone the heteromultimers of the serotonin-gated anion channel then proceed, as described above.

## II. Electrophysiological screening of endogenous serotonin-gated currents

If the above expression-cloning strategy is not successful, or in the alternative, one can identify MOD-1 like serotonin-gated responses in mammalian cells. Both acutely dissociated neurons and brain-derived cell lines may be screened. Once a cell type with a robust serotonin-gated, ionotropic, anion-selective current is identified, these cells are used as an enriched source from which to isolate mRNA coding for a MOD-1-like receptor. Poly(A)<sup>+</sup> RNA is then prepared from the positive tissue or cell line, and the same strategy as the one outlined in I is used to identify the clone(s) responsible for the response.

## III. Other approaches

In addition to the expression cloning strategies outlined above in sections I and II, complementary approaches can be taken to identify mammalian serotonin-gated chloride channels.

094240"2296550

A. EST databases are be systematically combed for sequences with similarities to the *mod-1* cDNA or protein sequence. This process is greatly enhanced by the identification of regions of the MOD-1 protein that allow it to be gated specifically by serotonin, and the regions of the protein that are predicted to be important for allowing anions to pass freely through the channel pore. Existing search algorithms for transmembrane topology (MEMSAT, TMAP, PHDtopology), protein fold motifs (TOPITS, UCLA-DOE Structure Prediction Server), and three-dimensional structures (SCOP) are used to search for sequences that may be clear candidates for mammalian homologs of MOD-1. Full-length cDNAs for such candidates are obtained and tested in the above-described oocyte expression system for the desired response.

B. Hybridization techniques are used to clone additional serotonin-gated anion channels. These techniques are well known to those skilled in the art, and are described, for example, in Ausubel et al., *Current Protocols in Molecular Biology*, John Wiley & Sons, New York, NY, 1990, and *Guide to Molecular Cloning Techniques*, 1987, S.L. Berger and A.R. Kimmel, eds., Academic Press, New York. If desired, a combination of different oligonucleotide probes may be used for the screening of a recombinant DNA library. The oligonucleotides are, for example, labeled with <sup>32</sup>P using methods known in the art, and the detectably-labeled oligonucleotides are used to probe filter replicas from a recombinant DNA library. Recombinant DNA libraries (for example, human cDNA libraries, such as brain or testes-derived cDNA libraries) are prepared according to methods well known in the art, for example, as described in Ausubel et al., *supra*, or are obtained

from commercial sources.

For detection or isolation of closely related serotonin-gated anion channel sequences, high stringency hybridization conditions are employed; such conditions include hybridization at about 42°C and about 50% formamide; a first wash at  
5 about 65°C, about 2X SSC, and 1% SDS; followed by a second wash at about 65°C and about 0.1% SDS, 1X SSC. Lower stringency conditions for detecting a gene encoding a serotonin-gated anion channel having less sequence identity to the nematode *mod-1* gene described herein include, for example, hybridization at about 42°C in the absence of formamide; a first wash at about 42°C, about 6X  
10 SSC, and about 1% SDS; and a second wash at about 50°C, about 6X SSC, and about 1% SDS.

C. Oligonucleotides that partially encode, or are complementary to nucleic acids encoding serotonin-gated anion channel-specific oligonucleotides are used as primers in PCR cloning strategies. Such PCR methods are well known in the art  
15 and are described, for example, in *PCR Technology*, H.A. Erlich, ed., Stockton Press, London, 1989; *PCR Protocols: A Guide to Methods and Applications*, M.A. Innis, D.H. Gelfand, J.J. Sninsky, and T.J. White, eds., Academic Press, Inc., New York, 1990; and Ausubel et al., *supra*. Again, sequences corresponding to sequences thought to encode amino acids important for serotonin-gated anion  
20 channel structure or function are preferred for use in isolating other sequences structurally and/or functionally related to a serotonin-gated anion channel. Such sequences are used to screen cDNA, as well as genomic DNA libraries. The sequences also include those that are not known to be important for serotonin-gated anion channel structure or function.



to treat disorders associated with serotonin-mediated cellular responses), or leads for such compounds, that can be used in human patients. In particular examples, compounds that specifically down-regulate or specifically increase serotonin-gated anion channel biological activity or the biological activity of their human  
5 homologs are considered useful in the invention. Also useful in the invention, are compounds that specifically affect other serotonin receptors and a serotonin-gated anion channel, or other serotonin receptors, but specifically not a serotonin-gated anion channel. In general, the screening methods of the invention involve screening any number of compounds for therapeutically active agents by  
10 employing any number of *in vitro* or *in vivo* experimental systems. Exemplary methods useful for the identification of such compounds are detailed below.

The methods of the invention simplify the evaluation, identification, and development of active agents for the treatment of conditions associated with serotonin-mediated cellular responses such as depression, panic disorders,  
15 obsessive compulsive disorder, sleep disorders, eating disorders, nausea and vomiting, other gastrointestinal disorders, and migraines, and the side-effects associated with these drugs. In general, the screening methods provide a facile means for selecting natural or synthetic product extracts or compounds of interest from a large population. These candidates are further evaluated and condensed to  
20 a few active and selective materials. Constituents of this pool are then purified and evaluated using methods described herein, to determine their ability to modulate serotonin-mediated responses and conditions.

Below we describe screening methods for evaluating the efficacy of a compound for use in the treatment of diseases associated with serotonin-mediated  
25 neurotransmission.

### Test Extracts and Compounds

In general, novel drugs for the treatment of serotonin-mediated cellular responses and conditions are identified from large libraries of both natural product, or synthetic (or semi-synthetic) extracts or chemical libraries according to methods known in the art. Those skilled in the field of drug discovery and development will understand that the precise source of test extracts or compounds is not critical to the screening procedure(s) of the invention. Accordingly, virtually any number of chemical extracts or compounds can be screened using the exemplary methods described herein. Examples of such extracts or compounds include, but are not limited to, plant-, fungal-, prokaryotic- or animal-based extracts, fermentation broths, and synthetic compounds, as well as modifications of existing compounds. Numerous methods are also available for generating random or directed synthesis (e.g., semi-synthesis or total synthesis) of any number of chemical compounds, including, but not limited to, saccharide-, lipid-, peptide-, and nucleic acid-based compounds. Synthetic compound libraries are commercially available from Brandon Associates (Merrimack, NH) and Aldrich Chemical (Milwaukee, WI). Alternatively, libraries of natural compounds in the form of bacterial, fungal, plant, and animal extracts are commercially available from a number of sources, including, but not limited to, Biotics (Sussex, UK), Xenova (Slough, UK), Harbor Branch Oceanographics Institute (Ft. Pierce, FL), and PharmaMar, U.S.A. (Cambridge, MA). In addition, natural and synthetically produced libraries are produced, if desired, according to methods known in the art, e.g., by combinatorial-chemistry methods or standard extraction and fractionation methods. Furthermore, if desired, any library or compound is readily modified



using standard chemical, physical, or biochemical methods.

In addition, those skilled in the art of drug discovery and development readily understand that methods for dereplication (e.g., taxonomic dereplication, biological dereplication, and chemical dereplication, or any combination thereof) or the elimination of replicates or repeats of materials already known for their effects on compounds associated with serotonin-mediated cellular responses should be employed whenever possible.

When a crude extract is found to affect serotonin-mediated responses or conditions, further fractionation of the positive lead extract is necessary to isolate chemical constituents responsible for the observed effect. Thus, the goal of the extraction, fractionation, and purification process is the careful characterization and identification of a chemical entity within the crude extract having activities that affect serotonin-mediated cellular responses. The same *in vivo* and *in vitro* assays described herein for the detection of activities in mixtures of compounds can be used to purify the active component and to test derivatives thereof. Methods of fractionation and purification of such heterogenous extracts are known in the art. If desired, compounds shown to be useful agents for the treatment of pathogenicity are chemically modified according to methods known in the art. Compounds identified as being of therapeutic value are subsequently analyzed using any standard animal model of serotonin-mediated cellular responses known in the art.

There now follow examples of systems useful for evaluating the efficacy of a molecule or compound in treating (or preventing) a condition associated with serotonin-mediated cellular responses, and the side-effects resulting from the treatment of these conditions.

Assays to be used for identifying compounds that affect serotonin-mediated cellular responses include assaying locomotion rates of nematodes exposed to test compounds. It also includes adding a test compound to a cell and assaying serotonin-gated anion channel expression at the nucleic acid level or at the polypeptide level. The changes in serotonin-gated anion channel RNA levels can be monitored by Northern blots, or by highly sensitive quantitative RT-PCR assays. The changes in the levels of serotonin-gated anion channel polypeptide can be monitored through the use of antibodies, including standard Western blot analyses, and immunohistochemistry.

The invention also includes assays that measure ionic current through a channel as a means of identifying a compound that affects serotonin-mediated cellular responses. In the presence of an agonist, the serotonin-gated anion channel is likely to be activated, and this will lead to an increase in the current carried through the channel. This change in current flow can be measured using standard electrophysiological methods. In the presence of an antagonist, the serotonin-gated anion channel is likely to be refractory to the application of serotonin, and less, or no current will pass through the channel. Such changes can be measured using standard electrophysiological methods.

The invention also includes assays that measure the concentration of serotonin, or a test compound required to activate a serotonin-gated anion channel, as compared to the 5-HT<sub>3</sub> receptor. In both HEK cell and *Xenopus* oocyte expression systems, serotonin-gated currents are detectable at lower concentration of agonist (10-50 nM) for MOD-1 receptors than for 5-HT<sub>3</sub> receptors(>10  $\mu$ M). This observation provides a tool for exploring the basis of dose-dependent clinical responses. Effects produced by low-dose serotonin or test compounds may be

mediated by serotonin-gated anion channels, while effects at 100-fold higher serotonin or test compound concentrations are likely to arise from a combination of serotonin-gated anion channels and 5-HT<sub>3</sub> type receptor activation.

#### Example 8: Diagnostic probes

5 cDNA fragments can be used as hybridization probes for allelic markers for haplotype analysis of human disorders or conditions linked to the locus of a serotonin-gated anion channel locus. Such analyses can also be performed using other standard techniques, such as PCR.

#### Example 9: Pharmacogenetics of responses to therapeutics for serotonin-mediated conditions

10 A serotonin-gated anion channel can serve as a marker for determining how an individual might respond to a given therapeutic for a serotonin-mediated condition. Genetic analysis of an individual's serotonin-gated anion channel locus can be completed as described above. Once a group of major mutant alleles has  
15 been established, PCR based genotyping assays can be developed to make a molecular diagnosis of an abnormal serotonin-gated anion channel. When information regarding how individuals with a specific serotonin-gated anion channel allele respond to particular therapies is combined with molecular diagnosis techniques, it is feasible to select an optimal therapy to treat serotonin-mediated  
20 conditions.

Example 10: Therapies that modulate conditions mediated by serotonergic pathways: Effect of serotonin-gated anion channel genotype

Patients experiencing a specific condition, or set of conditions, mediated by serotonin are placed into one of four treatment groups: placebo or 3 increasing doses of a given drug. During and at the end of the treatment period, the patients are evaluated for the effect of the drug on modulation of the condition(s). The patients are also evaluated for side-effects experienced as a result of drug administration. Patients are also characterized for the presence of various alleles of a serotonin-gated anion channel, and optionally, other serotonin-mediated receptor alleles and/or their respective protein levels. Results of these studies can be correlated to design therapies that provide optimal relief from the serotonin-mediated condition(s), with the fewest side-effects, in a patient with a specific serotonin-gated anion channel allele, or particular protein levels. In addition, other known serotonin-mediated receptors may be characterized and included in the development of the therapeutic protocol.

The patients can also be genotyped for polymorphisms in specific enzymes required for the metabolism of a given therapeutic used in the treatment of serotonin-mediated conditions. Such enzymes to be assessed include thiopurine S-methyltransferase (TPMT), dihydropyrimidine dehydrogenase (DPD), aldehyde dehydrogenase (ALDH), glutathione S-transferase (GST), uridine diphosphate glucuronosyl-transferase (UGTs), and cytochrome P450 enzymes. Once knowledge of an individual's drug metabolism profile is obtained, therapies to treat serotonin-mediated conditions can be more optimally designed to provide maximal efficacy with minimal side-effects.

## Antisense RNA

1. *Design of antisense systems.* One way in which antisense RNA can be synthesized is through a system. One example of a system includes, but is not limited to a complete panel of adenovirus constructs. The panel may consist of approximately four types of recombinant virus: A) A sense orientation virus for each serotonin-gated anion channel open reading frame; these viruses are designed to massively overexpress the recombinant protein in infected cells. B) Antisense orientation viruses in which the viral promoter drives the synthesis of an mRNA of opposite polarity to the serotonin-gated anion channel RNA, thereby shutting off host cell synthesis of the targeted protein coding region. C) Sub-domain expression viruses; these constructs express only a partial serotonin-gated anion channel protein in infected cells. D) Control viruses; functional analysis of serotonin-gated anion channel requires suitable positive and negative controls for comparison.
2. *Confirmation of recombinant adenovirus function.* Verification of the sense adenovirus function involves infection of tissue culture cells and determination of protein expression levels. This can be done by Western blot analysis. Functional analysis of the antisense viruses may be done at the RNA level using either Northern blots of total RNA harvested from infected tissue culture cells or ribonuclease protection assays. Western blot analysis of infected cells may be used to determine whether the expressed antisense RNA interferes with serotonin-gated anion channel expression in the host cell.

3. *Documentation that serotonin-gated anion channel over-expression results in increased serotonin-gated anion channel activity.* Determination of whether serotonin-gated anion channel over-expression results in increased serotonin-gated channel activity can be done by measuring the ionic current across a membrane elicited by serotonin using standard voltage-clamping techniques. The surface area of the membrane to be analyzed is pre-determined and remains constant for assaying both over-expressing and control samples.

4. *Documentation that antisense serotonin-gated anion channel over-expression results in decreased serotonin-gated anion channel activity.* Having confirmed that serotonin-gated anion channel over-expression renders cells more likely to exhibit increased serotonin-gated anion channel activity, one may examine whether the antisense adenoviruses render the same cells resistant to the channel activity, using the above-described methods.

5. *Identification of antisense oligonucleotides.* Concomitant to the adenovirus work, a series of antisense oligonucleotides to various regions of a serotonin-gated anion channel can also be designed. A generally-accepted model of how antisense oligonucleotides function proposes that the formation of RNA/DNA duplexes in the nucleus activates cellular RNase H enzymes which enzymatically degrade the mRNA component of the hybrid. Virtually any region of the mRNA can be targeted, and therefore choosing an appropriate sequence to target is somewhat empirical. One site frequently targeted is the translation initiation site. Alternatively, one can design antisense oligonucleotides that systematically “walk” down a nucleic acid sequence of interest. Many factors, including secondary

structure of the target mRNA and the binding affinity of the targeted sequence determine whether a particular oligonucleotide will be effective, necessitating several oligos for a serotonin-gated anion channel. Oligonucleotides to serotonin-gated anion channel mRNA can be made based on the available computer  
5 algorithms for predicting binding affinities and mRNA secondary structures. These and other oligonucleotides may be tested for their ability to target their respective mRNAs for degradation using Northern blot analysis.

6. *Optimization of oligonucleotides.* A secondary round of oligonucleotides may be made when effective target regions have been identified. These oligonucleotides  
10 target sequences in the immediate vicinity of the most active antisense oligonucleotides identified using methods such as those provided above. A second round of testing by Northern blot analysis may be required.

7. *Testing antisense oligonucleotides in vitro.* Following successful identification and optimization of targeting oligonucleotides, one may test these antisense  
15 oligonucleotides in tissue culture cells. Experimental procedures may parallel those used in the recombinant antisense adenovirus work. Negative control oligonucleotides with mismatch sequences are used to establish baseline or non-specific effects. Assisted transfection of the oligonucleotides using, for example, cation lipid carriers may be compared to unassisted transfection. Confirmation of  
20 the effectiveness of specific antisense oligonucleotides prompts synthesis of oligonucleotides with modified phosphodiester linkages, such as phosphorothioate or methylimino substituted oligos. These modified oligonucleotides may also be tested *in vitro*.

Another therapeutic approach within this invention involves administration of recombinant protein fragments or antibodies to a serotonin-gated anion channel, either directly to the site where modulation of serotonin transmission is desirable (for example, by injection) or by systemic administration (for example, by any conventional recombinant protein administration technique).

The dosage of serotonin-gated anion channel, the serotonin-gated anion channel fragment, serotonin-gated anion channel mutant protein, or antibody to a serotonin-gated anion channel depends on a number of factors, including the size and health of the individual patient, but, generally, between 0.1 mg and 100 mg inclusive are administered per day to an adult in any pharmaceutically acceptable formulation.

#### Administration

A serotonin-gated anion channel mutant protein or protein fragment, gene encoding the same, gene encoding a serotonin-gated anion channel antisense RNA, or modulator of a serotonin-gated anion channel may be administered within a pharmaceutically-acceptable diluent, carrier, or excipient, in unit dosage form. Conventional pharmaceutical practice may be employed to provide suitable formulations or compositions to administer the compounds to patients suffering from a disease or condition associated with the serotonergic pathway.

Administration may begin before the patient is symptomatic. Any appropriate route of administration may be employed, for example, administration may be parenteral, intravenous, intraarterial, subcutaneous, intramuscular, intracranial, intraorbital, ophthalmic, intraventricular, intracapsular, intraspinal, intracisternal, intraperitoneal, intranasal, aerosol, suppository, or oral administration. For example, therapeutic



formulations may be in the form of liquid solutions or suspensions; for oral administration, formulations may be in the form of tablets or capsules; and for intranasal formulations, in the form of powders, nasal drops, or aerosols.

Methods well known in the art for making formulations are found, for example, in "Remington's Pharmaceutical Sciences," (Remington: The Science and Practice of Pharmacy, 19<sup>th</sup> ed., A.R. Gennaro, ed., Mack Publishing Co., Easton, PA, 1995). Formulations for parenteral administration may, for example, contain excipients, sterile water, or saline, polyalkylene glycols such as polyethylene glycol, oils of vegetable origin, or hydrogenated naphthalenes. Biocompatible, biodegradable lactide polymer, lactide/glycolide copolymer, or polyoxyethylene-polyoxypropylene copolymers may be used to control the release of the compounds. Other potentially useful parenteral delivery systems for serotonin-gated anion channel modulatory compounds include ethylene-vinyl acetate copolymer particles, osmotic pumps, implantable infusion systems, and liposomes. Formulations for inhalation may contain excipients, for example, lactose, or may be aqueous solutions containing, for example, polyoxyethylene-9-lauryl ether, glycocholate and deoxycholate, or may be oily solutions for administration in the form of nasal drops, or as a gel.

If desired, treatment with serotonin-gated anion channel mutant proteins or serotonin-gated anion channel fragments, related genes, or other modulatory compounds may be combined with more traditional therapies for therapies associated with serotonin-mediated cellular responses.

### Example 11: Transgenic Mice

Transgenic mouse expression vectors, including neuronal, testes, and smooth muscle cell-specific promoter constructs can be constructed. Founder mice that are viable for most of these constructs can be identified, and breeding colonies can be developed. These mice will likely be prone to modulation of the serotonin-gated anion channel within tissue types in which the promoter is active. Thus the mice provide an excellent resource for testing the efficacy of antisense oligonucleotides, and for screening for therapeutics associated with serotonin-mediated cellular responses and the side-effects associated with these therapeutics. Standard mouse drug screening models and gene delivery protocols may be employed to utilize the mice for this purpose.

What is claimed is:

1. A substantially pure nucleic acid sequence encoding a serotonin-gated anion channel.

2. A substantially pure polypeptide, said polypeptide being a serotonin-gated anion channel.

5           3. The nucleic acid sequence of claim 1, wherein said serotonin-gated anion channel is a chloride channel.

4. The polypeptide sequence of claim 2, wherein said serotonin-gated anion channel is a chloride channel.

10           5. The nucleic acid sequence of claim 1, wherein said serotonin-gated anion channel is MOD-1.

6. The polypeptide sequence of claim 2, wherein said serotonin-gated anion channel is MOD-1.

7. An antibody that specifically binds to a serotonin-gated anion channel.

8. A *Caenorhabditis elegans* (*C. elegans*) strain having a mutant *mod-1* gene.

9. A method for identifying a compound that modulates the biological activity of a serotonin-gated anion channel, said method comprising the steps of:

(a) administering a test compound to a serotonin-gated anion channel;  
and

5 (b) assaying a modulation in the biological activity of said serotonin-gated anion channel.

10 10. A diagnostic probe for detecting conditions associated with a serotonin-mediated cellular response, said probe comprising a means for measurement of a serotonin-gated anion channel.

10 11. A method for characterizing a drug associated with serotonin-mediated cellular response, said method comprising detecting a modulation in the activity of a serotonin-gated anion channel when said channel is exposed to said drug.

15 12. A method for decreasing serotonin-gated anion channel function, said method comprising administering an antisense RNA that decreases the level of a serotonin-gated anion polypeptide.

13. A method for decreasing serotonin-gated anion channel function, said method comprising administering an antibody that binds to a serotonin-gated anion channel polypeptide.

14. A method for modulating serotonin-gated anion channel function, said method comprising administering a nucleic acid vector encoding a serotonin-gated anion channel, said administering being sufficient to modulate serotonin-gated anion channel activity.

5           15. A method for identifying a gene that is structurally related to a gene encoding a serotonin-gated anion channel, said method comprising identifying a gene with a probe derived from said serotonin-gated anion channel gene or a product encoded by said serotonin-gated anion channel gene.

16. An isolated gene identified by the method of claim 85.

10           17. A transgenic animal that over-expresses a serotonin-gated anion channel.

18. A transgenic animal that under-expresses a serotonin-gated anion channel.

19. A transgenic animal that expresses a dominant negative serotonin-gated anion channel.

20. A method for identifying a compound that modulates the activity of a serotonin-gated anion channel, said method comprising the steps of:

- (a) exposing a nematode to a test compound;
- (b) assaying the locomotion rate of said nematode; and
- 5 (c) comparing said locomotion rate to that of a control nematode receiving no test compound, wherein a modulation in said locomotion rate indicates a compound that modulates the activity of a serotonin-gated anion channel.

21. A method for identifying a compound that modulates the activity of a serotonin-gated anion channel in a liquid locomotion assay, said method comprising the steps of:

- (a) exposing a nematode to a test compound;
- (b) quantifying the number of nematodes actively swimming after exposure to said test compound; and
- 15 (c) comparing the number of said actively swimming nematodes to that of control nematodes receiving no test compound, wherein a modulation in said number of actively swimming nematodes indicates a compound that modulates the activity of a serotonin-gated anion channel.







```

1150      1160      1170      1180      1190      1200
TGAACACGAT CATGAGCAAC TACACGAAAA TGTTGCCCGA CGCGGAGGAC AGCGTACAAG

1210      1220      1230      1240      1250      1260
TTAATATTGA GATTCATGTA CAGGTTGGTA GACTCTATAA TTGCACACCA ATATGTGAAA

1270      1280      1290      1300      1310      1320
GTTTTCTTTA AAATTAAACT GCTGTAAATG ACTTTTGAAT AAGTTTATCA GATAGAAATT

1330      1340      1350      1360      1370      1380
GTCTGAACTT TTCGATTCAA ACTTCCGAA CTTCAAAGCG GTTCCAAATT ACTCACTTCC

1390      1400      1410      1420      1430      1440
ATTTATCTCT TTGCTACAAT TTCTCCCACA AAGCCTTTTT CTTCAATTAA CGTTCTTTTT

1450      1460      1470      1480      1490      1500
TATGTCGTTG TTCTTACAAA CAATTCGTC TCCTTGATGA ACTGCTTGAA CTGAGAATAG

1510      1520      1530      1540      1550      1560
TCACATGAGG ATAAATTTGA TGAATGACA AGTTTTGTGC CCAGAAGGCA GTTTTGCCT

1570      1580      1590      1600      1610      1620
GAACTTGTTT AGTTGCAGAC ACATCTCAAA ACACAGAAGA TGAGTGGAAA ACTAGTGAGA

1630      1640      1650      1660      1670      1680
GACTGCCAAA AGTCGAAGGG ATAATGAAAA TTTGTTGCAA ATGAATTCTG CGAAGTTATG

1690      1700      1710      1720      1730      1740
TGAAAAATTA TTGGATTGGG AGTTGTGGGA GTGAAGAGAT GGGTCAAAAG CCATCAATCT

1750      1760      1770      1780      1790      1800
TGAATGCTTC GGTCAAAGAT TTGTTTCTCA TATGTTTACA AACTGAAAA CAATCTATCC

1810      1820      1830      1840      1850      1860
TAGAAATGTT TGAACCACCC TCTAAAGTCC TTCCGTATAT TTTTTCATCT TTATACCGAC

1870      1880      1890      1900      1910      1920
CAGAATTCAA GAGTTGTTTG AAATAACTTC CTCTTTTTTG GAGAATATGT ACTCAGATT

1930      1940      1950      1960      1970      1980
TTACATTCAA AATTTATATA TTTTCAAATA GAAAAAGTGC CAAGTACCAG AAACTTTTAT

1990      2000      2010      2020      2030      2040
CAAGTTGGCG GCACTTTGGA GAGTGAATTT GATGAAAAAG TGTTTGATAA GTTTGTCGGG

2050      2060      2070      2080      2090      2100
CAAACGGTTC CCCTGGGTGG GGAAATGGTG GCATTTTTTG AAACATTTTC ATAGTCGAAG

2110      2120      2130      2140      2150      2160
AAGTGAACA AGAAATTGG AAAATAGAGA TACATATGTA TATGAAAATA GAATTGAACA

2170      2180      2190      2200      2210      2220
GGAACCTATT TTTATTTTCA GGATATGGGA AGCTTGAATG AAATATCATC CGACTTTGAA

2230      2240      2250      2260      2270      2280
ATTGACATTT TATTCCTCA ACTGTGGCAT GACTCGGCAC TTTCTTTTGC TCATCTCCG

```

Fig. 1  
Page 2

2290	2300	2310	2320	2330	2340
GCTTGTAAGC	GGTAAGAAAT	CTTTGTATTA	GAAGGGAAAA	ATATTTAAAT	TAATGAAATT
2350	2360	2370	2380	2390	2400
TCAGAAATAT	CACAAATGGAA	ACACGACTTT	TACCTAAGAT	TTGGTCTCCA	AACACGTGTA
2410	2420	2430	2440	2450	2460
TGATTAATTC	AAAACGAACA	ACCGTCCATG	CATCACCATC	GGAAAATGTG	ATGGTTATTC
2470	2480	2490	2500	2510	2520
TGTACGAGGT	ATGATTTTTG	ATTTTGTGAC	GTCACAAACA	GAGCATGTCT	AAGGGCATGT
2530	2540	2550	2560	2570	2580
TGTAGCAAGA	AAAAACGGA	TTCTTGCTC	TGTCGACGTT	TCCTAAGTAT	TGTGAATTAT
2590	2600	2610	2620	2630	2640
TTATAATACA	TCACTCTAAT	TACGTGAATA	CTTACACCTT	TAAGTGGGTG	AAGGATAAAA
2650	2660	2670	2680	2690	2700
TAGAGAAGGA	GACGTTGAAA	AAGCTCTTCG	GTAGATTAAA	GAGTCTAGAA	TCGACATATG
2710	2720	2730	2740	2750	2760
TATTCATGTT	TCTCGGTTCA	GGGAAATAAG	TGATTTTGGC	GAAAAAGAGT	TAGACGACAT
2770	2780	2790	2800	2810	2820
TTTTTTAGAA	ACTAAACTA	TATTCTCGAA	CCCAAATCAG	TCTAATGGTT	TTCAGCAAAA
2830	2840	2850	2860	2870	2880
AGTATGAAAT	ATACAATGTT	TGTTTCAGAA	TACCCAGTAC	AAAATTTGAA	GTTTTTTCAGA
2890	2900	2910	2920	2930	2940
ATGGAACAGT	CTGGATTAACT	CATCGTCTTA	GTGTCAAATC	ACCTTGCAAT	TTGGATCTGC
2950	2960	2970	2980	2990	3000
GACAGTTTCC	TTTCGATACT	CAAACCTGCA	TATTAATCTT	TGAATCCTAT	AGTCATAACT
3010	3020	3030	3040	3050	3060
CAGAAGAAGT	TGAACTTCAT	TGGATGGAAG	AAGCTGTCAC	ATTAATGAAG	CCAATTCAAC
3070	3080	3090	3100	3110	3120
TTCTTGACTT	TGATATGGTT	CATTATTCAA	CTAAAAAGGA	AACTTTACTC	TATCCAAACG
3130	3140	3150	3160	3170	3180
GGTACTGGGA	TCAGCTTCAA	GTTACTTTCA	CTTTCAAACG	ACGATATGGA	TTCTATATTA
3190	3200	3210	3220	3230	3240
TTCAAGCCTA	TGTTCCAACA	TATCTTACAA	TCATTGTATC	TTGGGTTTCA	TTCTGCATGG
3250	3260	3270	3280	3290	3300
AACCAAAAGC	TCTGCCGGCA	AGAACAACCTG	TCGGAATCTC	ATCTCTTCTA	GCTCTTACTT
3310	3320	3330	3340	3350	3360
TCCAGTTTGG	AAATATTTTG	AAAAATCTTC	CAAGGGTTTC	ATATGTGAAA	GGTTTGTTTT
3370	3380	3390	3400	3410	3420
TTTTCTTTTT	CAAAACAAATA	AAAAAAAAGA	TAAACAAATA	TTTGTTTCAG	CAATGGATGT



4570            4580            4590            4600            4610            4620  
 CACACACACA CACACATTCA CGAAACTTTG TGTGTGTTAT GTTACTTATA TGTATCTTT

4630            4640            4650            4660            4670            4680  
 TCTGTCTGAT CATGGTTTTC GGAAGTATA TGTGTTAATC GGAAGTATA TGTGAGCCAC

4690            4700            4710            4720            4730            4740  
 ATTGATTAAA CCTGTGAGAG ATGCCCATTG GTACTCATTG TACGACTGTC TCATGTCCAA

4750            4760            4770            4780            4790            4800  
 ACACCATGTT TATTGTAATT ACCAGGCTAC TATTTGCAGA TGCGATCAAC ATCACCACCT

4810            4820            4830            4840            4850            4860  
 CCACCATCTG GATGTCTGGC CAGATTCAT CCGGAAGCAG TGGACAAATT CTCCATTGTA

4870            4880            4890            4900            4910            4920  
 GCTTTTCCAT TGGCATTAC AATGTTAAT GTTAGTTAAT CCACAGTTAA AAATTTCCAT

4930            4940            4950            4960            4970            4980  
 AATCATAAAT ATCTCGACTT TTCAGCTTGT CTAAGTGGTG CACTATTTGT CTCAAATTT

4990            5000            5010            5020            5030            5040  
 CGATCAAAC TATCAGTAT TGAAGTTTAT CCCTTTTAA TCCAATAATT CACAGTTGCC

5050            5060            5070            5080            5090            5100  
 GGTATCTACC TCCATTCTTT TCCGATGATT CGCAGTTTTT CACAGGGTTC AAATGTATCT

5110            5120            5130            5140            5150            5160  
 CGTTCAATCT TTTTATGGTT ATTTCTCTTG AATGTCCATT TTAATATTTA TAGAACACTT

5170            5180            5190            5200            5210            5220  
 TTATGTACAT TGTGTTGGTA TTCAATTCGA AAAACAATGA AATTTATTTT TAAATAACTG

5230            5240            5250            5260            5270            5280  
 CGTTTCTGGG GTTTCTATCA GCACTTACTA GCTGACAAAA ACTTTTCCGT ATTCGGAATT

5290            5300            5310            5320            5330            5340  
 AGATTTTAT GCAAGCAATG TTTCATTTTT ACACAGTATA GTATTTATTC TTACTTTTGA

5350            5360            5370            5380            5390            5400  
 TTATATTGCT CGCACCTAA ATGACAGGTA TTAGAAATTA ACCGCTTTTC AGAGTATTTT

5410            5420            5430            5440            5450            5460  
 TAATCTTCTT AGTACTAGTT TAGTTCTTTA AATAAGAAAC CATCTAGTTT TTCATTATCA

5470            5480            5490            5500            5510            5520  
 CTCAACTTCA GTCGGACAAA TTTTAAATTT TTTACTCGAT AAAAAATTT TATAATTCAG

5530            5540            5550  
 ACAAATTATG TCTTCTCATT TTTGATCGCT

10 20 30 40 50  
 ATGAAGTTTA TTCCTGAAAT CACACTACTC TTGCTTTTAT TTGTACACTC  
 60 70 80 90 100  
 TACACAGGCT AAAGGAAAAC GACGGAAATG TCCGGAGGGT GCGTGGTCGG  
 110 120 130 140 150  
 AAGGAAAGAT TATGAACACG ATCATGAGCA ACTACACGAA AATGTTGCCC  
 160 170 180 190 200  
 GACGCGGAGG ACAGCGTACA AGTTAATATT GAGATTCATG TACAGGATAT  
 210 220 230 240 250  
 GGGAAGCTTG AATGAAATAT CATCCGACTT TGAAATTGAC ATTTTATTCA  
 260 270 280 290 300  
 CTCAACTGTG GCATGACTCG GCACTTTCTT TTGCTCATCT TCCGGCTTGT  
 310 320 330 340 350  
 AAGCGAAATA TCACAATGGA AACACGACTT TTACCTAAGA TTTGGTCTCC  
 360 370 380 390 400  
 AAACACGTGT ATGATTAATT CAAAACGAAC AACCGTCCAT GCATCACCAT  
 410 420 430 440 450  
 CGGAAAATGT GATGGTTATT CTGTACGAGA ATGGAACAGT CTGGATTAAAC  
 460 470 480 490 500  
 CATCGTCTTA GTGTCAAATC ACCTTGCAAT TTGGATCTGC GACAGTTTCC  
 510 520 530 540 550  
 TTTGATACT CAAACTTGCA TATTAATCTT TGAATCCTAT AGTCATAACT  
 560 570 580 590 600  
 CAGAAGAAGT TGAACCTCAT TGGATGGAAG AAGCTGTCAC ATTAATGAAG  
 610 620 630 640 650  
 CCAATTCAAC TTCCTGACTT TGATATGGTT CATTATTCAA CTAAAAAGGA  
 660 670 680 690 700  
 AACTTTACTC TATCCAAACG GGTACTGGGA TCAGCTTCAA GTTACTTTCA  
 710 720 730 740 750  
 CTTTCAAACG ACGATATGGA TTCTATATTA TTCAAGCCTA TGTTCCAACA  
 760 770 780 790 800  
 TATCTTACAA TCATTGTATC TTGGGTTTCA TTCTGCATGG AACCAAAAGC  
 810 820 830 840 850  
 TCTGCCGGCA AGAACAACTG TCGGAATCTC ATCTCTTCTA GCTCTTACTT  
 860 870 880 890 900  
 TCCAGTTTGG AAATATTTTG AAAAATCTTC CAAGGGTTTC ATATGTGAAA  
 910 920 930 940 950  
 GCAATGGATG TGTGGATGCT TGGATGCATA TCATTTGTCT TCGGAACCAT


	1980	1981	1982	1983	1984	1985	1986	1987	1988	1989	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015	2016	2017	2018	2019	2020	2021	2022	2023	2024	2025	2026	2027	2028	2029	2030	2031	2032	2033	2034	2035	2036	2037	2038	2039	2040	2041	2042	2043	2044	2045	2046	2047	2048	2049	2050	2051	2052	2053	2054	2055	2056	2057	2058	2059	2060	2061	2062	2063	2064	2065	2066	2067	2068	2069	2070	2071	2072	2073	2074	2075	2076	2077	2078	2079	2080	2081	2082	2083	2084	2085	2086	2087	2088	2089	2090	2091	2092	2093	2094	2095	2096	2097	2098	2099	2100	2101	2102	2103	2104	2105	2106	2107	2108	2109	2110	2111	2112	2113	2114	2115	2116	2117	2118	2119	2120	2121	2122	2123	2124	2125	2126	2127	2128	2129	2130	2131	2132	2133	2134	2135	2136	2137	2138	2139	2140	2141	2142	2143	2144	2145	2146	2147	2148	2149	2150	2151	2152	2153	2154	2155	2156	2157	2158	2159	2160	2161	2162	2163	2164	2165	2166	2167	2168	2169	2170	2171	2172	2173	2174	2175	2176	2177	2178	2179	2180	2181	2182	2183	2184	2185	2186	2187	2188	2189	2190	2191	2192	2193	2194	2195	2196	2197	2198	2199	2200	2201	2202	2203	2204	2205	2206	2207	2208	2209	2210	2211	2212	2213	2214	2215	2216	2217	2218	2219	2220	2221	2222	2223	2224	2225	2226	2227	2228	2229	2230	2231	2232	2233	2234	2235	2236	2237	2238	2239	2240	2241	2242	2243	2244	2245	2246	2247	2248	2249	2250	2251	2252	2253	2254	2255	2256	2257	2258	2259	2260	2261	2262	2263	2264	2265	2266	2267	2268	2269	2270	2271	2272	2273	2274	2275	2276	2277	2278	2279	2280	2281	2282	2283	2284	2285	2286	2287	2288	2289	2290	2291	2292	2293	2294	2295	2296	2297	2298	2299	2300	2301	2302	2303	2304	2305	2306	2307	2308	2309	2310	2311	2312	2313	2314	2315	2316	2317	2318	2319	2320	2321	2322	2323	2324	2325	2326	2327	2328	2329	2330	2331	2332	2333	2334	2335	2336	2337	2338	2339	2340	2341	2342	2343	2344	2345	2346	2347	2348	2349	2350	2351	2352	2353	2354	2355	2356	2357	2358	2359	2360	2361	2362	2363	2364	2365	2366	2367	2368	2369	2370	2371	2372	2373	2374	2375	2376	2377	2378	2379	2380	2381	2382	2383	2384	2385	2386	2387	2388	2389	2390	2391	2392	2393	2394	2395	2396	2397	2398	2399	2400	2401	2402	2403	2404	2405	2406	2407	2408	2409	2410	2411	2412	2413	2414	2415	2416	2417	2418	2419	2420	2421	2422	2423	2424	2425	2426	2427	2428	2429	2430	2431	2432	2
--	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	---

Fig. 2

10	20	30	40	50
MKFIPEITLL	LLLFVHSTQA	KGKRRKCPEG	AWSEGKIMNT	IMSNYTKMLP
60	70	80	90	100
DAEDSVQVNI	EIHVQDMGSL	NEISSDFEID	ILFTQLWHDS	ALSFAPLPAC
110	120	130	140	150
KRNITMETRL	LPKIWSPNTC	MINSKRRTVH	ASPSENVMI	LYENGTWIN
160	170	180	190	200
HRLSVKSPCN	LDLRQFPFDT	QTCILIFESY	SHNSEEVELH	WMEEAVTLMK
210	220	230	240	250
PIQLPDFDMV	HYSTKKTLL	YPNGYWDQLQ	VTFTFKRRYG	FYIIQAYVPT
260	270	280	290	300
YLTIIWSWS	FCMEPKALPA	RTTVGISSLL	ALTFQFGNIL	KNLPRVSIVK
310	320	330	340	350
AMDVWMLGCI	SFVFGTMVEL	AFVCYISRCQ	NSVRNAERRR	ERMRNSQVWA
360	370	380	390	400
NGSCRTRSNG	YANGGSVISH	YHPTSNGNGN	NNRHDTQPVT	GRGSLHRNGP
410	420	430	440	450
PSPLNLQMTT	FDSEIPLTFD	QLPVSMESDR	PLIEEMRSTS	PPPPSGCLAR
460	470	480		
FHPEAVDKFS	IVAFPLAFTM	FNLVYWWHYL	SQTFDQNYQ	

Fig. 3

2





# ok103 is a 4135 bp deletion allele of *mod-1*




Fig. 5

```

      10      20      30      40      50      60
TCATGTTTCA CGGAACGACG AATTTATCCC GTCGTTTCTT CCTTTCCGTT TTAACTCATA

      70      80      90     100     110     120
TCTCTTCCTG GATCCTTCAG AGCTCTTGTC AATTCCTCAC GTTTTTTTTT GTTTTTTCGT

     130     140     150     160     170     180
CGTTTAATTG TGGAACACA TATCCGTCCT CTTTGAAACA GCATCAGAAA ACTTTCTGCT

     190     200     210     220     230     240
CTCCGTGTCC TTCTACTTAC TCTGATTGCC TTAGTTAGTC ACATCGCAAG CAACAATAA

     250     260     270     280     290     300
CTGCCAATGG GAGGAGCCAG TTGGAGCAGG GTGCGTGCTC GGTGCTCTTT TCAGAAGGTT

     310     320     330     340     350     360
TTCTCTTG TG CCAGCATGCT TTTTGTAGGC TGTGTCATCA CAATGAACAT GTGTGAGTTC

     370     380     390     400     410     420
ATCCGTCTGG ATTATTCTTT TTCTTACGTC TTCTGAGTAC TTCATACTTT CCAAATTTTT

     430     440     450     460     470     480
CAACTGAAC TTTCTTCTTT TCTCATTGAA GTGGTTTGGT TTTGGTCGCG TGATCAACGG

     490     500     510     520     530     540
ATCCTACTTT TTTGAAACAA AATGTTTTTG AAGTTTCACA GACTGATTTC GGGGTTTTTT

     550     560     570     580     590     600
CAAAGAATAT ATTCCCTCTC GAGCAAGAGA AAATTCCAGA AAATAGTAGT TTTTTTCAAT

     610     620     630     640     650     660
TAGTCGTTTC ATTTGTACTA GCTAAAAAC TTGCAACTTA TGGCTTTAAA ACATGTGTTG

     670     680     690     700     710     720
GCTTCATACA AAAACATTTA ACTAGTGTTT TTCCAGTTTT GTGTTCGTTT CATTTTCTCA

     730     740     750     760     770     780
CCAACTGAC AATAATTACT TTCTGTGAAC GTGTTTTGTA GGCAAGCTCC CGAATATTTT

     790     800     810     820     830     840
TTTCTCTTCT CACGTCTTGT TATTTTCTCG ATTTTATTTT CTGAATCTGT GCGGTTTTCA

     850     860     870     880     890     900
ATCAATTTGA TTGCGATAAT TATTCTATCA GAAATATATT TTCAGAAATC CAAATACTCC

     910     920     930     940     950     960
AGGTGCCAAT GCGGTGAAAG AAAATTATGA AGTTTATTCC TGAAATCACA CTA CTCTTGC

     970     980     990    1000    1010    1020
TTTTATTTGT AACTCTACA CAGGTTAGTT TCTCTTGAAT GTCCATTTTA ATATTTATAG

    1030    1040    1050    1060    1070    1080
AACACTTTTA TGTACATTGT GTTGGTATTC AATTCGAAAA ACAATGAAAT TTATTTCTAA

    1090    1100    1110    1120    1130    1140
ATAACTGCGT TTCTGGGGTT TCTATCAGCA CTTACTAGCT GACAAAAACT TTTCCGTATT

```

Fig. 6

1150	1160	1170	1180	1190	1200
CGGAATTAGA	TTTTTATGCA	AGCAATGTTT	CATTTTACAC	CAGTATAGTA	TTTATTCTTA
1210	1220	1230	1240	1250	1260
CTTTTGATTA	TATTGCTCGC	ACCCTAAATG	ACAGGTATTA	GAAATTAACC	GCTTTTCAGA
1270	1280	1290	1300	1310	1320
GTATTTTAA	TCTTCTTAGT	ACTAGTTTAG	TTCTTTAAAT	AAGAAACCAT	CTAGTTTTC
1330	1340	1350	1360	1370	1380
ATTATCACTC	AACTTCAGTC	GGACAAATTT	TAAATTTTTT	ACTCGATAAA	AAAATTTTAT
1390	1400	1410			
AAATTCAGACA	AATTATGTCT	TCTCATTTTT	GATCGCT		

10 20 30 40 50 60  
 TCATGTTTCA CGGAACGACG AATTTATCCC GTCGTTTCTT CCTTTCCGTT TTAAC TCATA  
 70 80 90 100 110 120  
 TCTCTTCCTG GATCCTTCAG AGCTCTTGTC AATTCCTCAC GTTTTTTTTT GTTTTTTCGT  
 130 140 150 160 170 180  
 CGTTTAATTG TGGAAACACA TATCCGTCCT CTTTGAAACA GCATCAGAAA ACTTTCTGCT  
 190 200 210 220 230 240  
 CTCCGTGTCC TTCTACTTAC TCTGATTGCC TTAGTTAGTC ACATCGCAAG CAACA ACTAA  
 250 260 270 280 290 300  
 CTGCCAATGG GAGGAGCCAG TTGGAGCAGG GTGCGTGCTC GGTGCTCTTT TCAGAAGGTT  
 310 320 330 340 350 360  
 TTCTCTTG TG CCAGCATGCT TTTT TGAGGC TGTGTCATCA CAATGAACAT GTGTGAGTTC  
 370 380 390 400 410 420  
 ATCCGTCTGG ATTATTCTTT TTCTTACGTC TTCTGAGTAC TTCATACTTT CCAAATTTT  
 430 440 450 460 470 480  
 CAACTGAAC TTTCTTCTTT TCTCATTGAA GTGGTTTGGT TTTGGTCGCG TGATCAACGG  
 490 500 510 520 530 540  
 ATCCTACTTT TTTGAAACAA AATGTTTTTG AAGTTTCACA GACTGATTTC GGGGTTTTTT  
 550 560 570 580 590 600  
 CAAAGAATAT ATTCCCTCTC GAGCAAGAGA AAATTC CAGA AAATAGTAGT TTTTTTCAAT  
 610 620 630 640 650 660  
 TAGTCGTTTC ATTTGTACTA GCTAAAAAC TTGCAACTTA TGGCTTTAAA ACATGTGTTG  
 670 680 690 700 710 720  
 GCTTCATACA AAAACATTTA ACTAGTGTTT TTCCAGTTTT GTGTTCGTTT CATTTTCTCA  
 730 740 750 760 770 780  
 CCAAAC TGAC AATAATTACT TTCTGTGAAC GTGTTTTGTA GGCAAGCTCC CGAATATTTT  
 790 800 810 820 830 840  
 TTTCTCTTCT CACGTCTTGT TATTTTCTCG ATTTTATTTT CTGAATCTGT GCGGTTTTCA  
 850 860 870 880 890 900  
 ATCAATTTGA TTGCGATAAT TATCTATCA GAAATATATT TTCAGAAATC CAAATACTCC  
 910 920 930 940 950 960  
 AGGTGCCAAT GCGGTGAAAG AAAATTATGA AGTTTATTCC TGAAATCACA CTACTCTTGC  
 970 980 990 1000 1010 1020  
 TTTTATTTGT AACTCTACA CAGGTAGTT GGTGATTCT AGATCTCTTG CCTCCTAGCT  
 1030 1040 1050 1060 1070 1080  
 TGCAAGGATA ATATAATTGA ATTGTTTTTG AGGAGTGCAA AGATTGAATA GTTTCTATA  
 1090 1100 1110 1120 1130 1140  
 TTTAGGCTAA AGGAAAACGA CGGAAATGTC CGGAGGGTGC GTGGTCGAA GGAAAGATTA

Fig. 7

1150	1160	1170	1180	1190	1200
TGAACACGAT	CATGAGCAAC	TACACGAAAA	TGTTGCCCCG	CGCGGAGGAC	AGCGTACAAG
1210	1220	1230	1240	1250	1260
TTAATATTGA	GATTCATGTA	CAGGTTGGTA	GA CTCTATAA	TTGCACACCA	ATATGTGAAA
1270	1280	1290	1300	1310	1320
GTTTTCTTTA	AAATTAAACT	GCTGTAAATG	ACTTTTGAAT	AAGTTTATCA	GATAGAAATT
1330	1340	1350	1360	1370	1380
GTCTGAACTT	TTCGATTCAA	ACTTTCCGAA	CTTCAAAGCG	GTTCCAAATT	ACTCACTTCC
1390	1400	1410	1420	1430	1440
ATTTATCTCT	TTGCTACAAT	TTCTCCCACA	AAGCCTTTTT	CTTCATTTAA	CGTTCTTTTT
1450	1460	1470	1480	1490	1500
TATGTCGTTG	TTCTTACAAA	CAATTTTCGTC	TCCTTGATGA	ACTGCTTGAA	CTGAGAATAG
1510	1520	1530	1540	1550	1560
TCACATGAGG	ATAAATTTGA	TGGAATGACA	AGTTTTGTGC	CCAGAAGGCA	GTTTTGCACT
1570	1580	1590	1600	1610	1620
GAAC TTGTTC	AGTTGCAGAC	ACATCTCAAA	ACACAGAAGA	TGAGTGGAAA	ACTAGTGAGA
1630	1640	1650	1660	1670	1680
GACTGCCAAA	AGTCGAAGGG	ATAATGAAAA	TTTGTGCAA	ATGAATTCTG	CGAAGTTATG
1690	1700	1710	1720	1730	1740
TGAAAAATTA	TTGGATTGGG	AGTTGTGGGA	GTGAAGAGAT	GGGTCAAAAAG	CCATCAATCT
1750	1760	1770	1780	1790	1800
TGAATGCTTC	GGTCAAAGAT	TTGTTTCTCA	TATGTTTACA	AACTGAAAA	CAATCTATCC
1810	1820	1830	1840	1850	1860
TAGAAATGTT	TGAACCACCC	TCTAAAGTCC	TTCCGTATAT	TTTTTCATCT	TTATACCGAC
1870	1880	1890	1900	1910	1920
CAGAATTCAA	GAGTTGTTTG	AAATAACTTC	CTCTTTTTTG	GAGAATATGT	ACTCAGATTT
1930	1940	1950	1960	1970	1980
TTACATTCAA	AATTTATATA	TTTTCAAATA	GAAAAAGTGC	CAAGTACCAG	AACTTTTAT
1990	2000	2010	2020	2030	2040
CAAGTTGGCG	GCACTTTGGA	GAGTGAATTT	GATGAAAAAG	TGTTTGATAA	GTTTGTCGGG
2050	2060	2070	2080	2090	2100
CAAAC TGGTC	CCCTGGGTGG	GGAAATGGTG	GCATTTTTTG	AAACATTTTC	ATAGTCGAAG
2110	2120	2130	2140	2150	2160
AAGTGAACA	AGAAAATTGG	AAAATAGAGA	TACATATGTA	TATGAAAATA	GAATTGAACA
2170	2180	2190	2200	2210	2220
GGAAC TTATT	TTTATTTTCA	GGATATGGGA	AGCTTGAATG	AAATATCATC	CGACTTTGAA
2230	2240	2250	2260	2270	2280
ATTGACATTT	TATTCAC TCA	ACTGTGGCAT	GA CTGCGCAC	TTTCTTTTGC	TCATCTTCCG

2290 2300 2310 2320 2330 2340  
 GCTTGTAAAGC GGTAAGAAAT CTTTGTATTA GAAGGGAAAA ATATTTAAAT TAATGAAATT  
 2350 2360 2370 2380 2390 2400  
 TCAGAAATAT CACAATGGAA ACACGACTTT TACCTAAGAT TTGGTCTCCA AACACGTGTA  
 2410 2420 2430 2440 2450 2460  
 TGATTAATTC AAAACGAACA ACCGTCCATG CATCACCATC GGAAAATGTG ATGGTTATTC  
 2470 2480 2490 2500 2510 2520  
 TGTACGAGGT ATGATTTTGTG ATTTTGTGAC GTCACAAACA GAGCATGTCT AAGGGCATGT  
 2530 2540 2550 2560 2570 2580  
 TGTAGCAAGA AAAAAACGGA TTCTTGTCTC TGTCGACGTT TCCTAAGTAT TGTGAATTAT  
 2590 2600 2610 2620 2630 2640  
 TTATAATACA TCACTCTAAT TACGTGAATA CTTACACCTT TAACTGGGTG AAGGATAAAA  
 2650 2660 2670 2680 2690 2700  
 TAGAGAAGGA GACGTTGAAA AAGCTCTTCG GTAGATTAAA GAGTCTAGAA TCGACATATG  
 2710 2720 2730 2740 2750 2760  
 TATTCATGTT TCTCGGTTCA GGGAAATAAG TGATTTTGGC GAAAAAGAGT TAGACGACAT  
 2770 2780 2790 2800 2810 2820  
 TTTTGTAGAAA ACTAAACTA TATTCTCGAA CCCAAATCAG TCTAATGGTT TTCAGCAAAA  
 2830 2840 2850 2860 2870 2880  
 AGTATGAAAT ATACAATGTT TGTTTCAGAA TACCCAGTAC AAAATTTGAA GTTTTTCAGA  
 2890 2900 2910 2920 2930 2940  
 ATGGAACAGT CTGGATTAAC CATCGTCTTA GTGTCAAATC ACCTTGCAAT TTGGATCTGC  
 2950 2960 2970 2980 2990 3000  
 GACAGTTTCC TTTCGATACT CAAACTTGCA TATTAATCTT TGAATCCTAT AGTCATAACT  
 3010 3020 3030 3040 3050 3060  
 CAGAAGAAGT TGAATTCAT TGGATGGAAG AAGCTGTCAC ATTAATGAAG CCAATTCAAC  
 3070 3080 3090 3100 3110 3120  
 TTCTGACTT TGATATGGTT CATTATTCAA CTAAAAAGGA AACTTTACTC TATCCAAACG  
 3130 3140 3150 3160 3170 3180  
 GGTACTGGGA TCAGCTTCAA GTTACTTTCA CTTTCAAACG ACGATATGGA TTCTATATTA  
 3190 3200 3210 3220 3230 3240  
 TTCAAGCCTA TGTTCCAACA TATCTTACAA TCATTGTATC TTGGGTTTCA TTCTGCATGG  
 3250 3260 3270 3280 3290 3300  
 AACCAAAAGC TCTGCCGGCA AGAACAACG TCGGAATCTC ATCTCTTCTA GTTCTTACTT  
 3310 3320 3330 3340 3350 3360  
 TCCAGTTTGG AAATATTTTG AAAAATCTTC CAAGGTTTC ATATGTGAAA GGTTTGTGTTT  
 3370 3380 3390 3400 3410 3420  
 TTTTCTTTTT CAACAAATA AAAAAAAGA TAAACAAATA TTTGTTTCAG CAATGGATGT

Fig. 7

3430 3440 3450 3460 3470 3480  
 GTGGATGCTT GGATGCATAT CATTTGTCTT CGGAACCATG GTAGAATTGG CATTTGTTTG  
 3490 3500 3510 3520 3530 3540  
 TTACATTTCC CGTTGTCAGA ACAGCGTAAG AAAGTGAGTT GGCATAAGAG TTTTCTCACG  
 3550 3560 3570 3580 3590 3600  
 TGGAGGGAAG TAATTAAATT TTGGGTGTCA TATGAAAATA TCAAAAACAA TATCAGGAAA  
 3610 3620 3630 3640 3650 3660  
 TTGAATTTCA CTATGATTTT GTAGTAAACA AATTACAGCG CGGAACGACG ACGGGAACGA  
 3670 3680 3690 3700 3710 3720  
 ATGAGAAATT CTCAGGTGTG GGCAAACGGA TCGTGTAGAA CTAGAAGCAA CGGGTATGCA  
 3730 3740 3750 3760 3770 3780  
 AACGGGGGAT CTGTAATCTC ACATTATCAT CCAACAAGCA ATGGAAATGG GAATAATAAT  
 3790 3800 3810 3820 3830 3840  
 CGACATGATA CACCTCAAGT TACTGGAAGG TTAGCAATCT CTATGATAGC ATTTATCAAT  
 3850 3860 3870 3880 3890 3900  
 TATTAAAGAA CTCTGGAATT AGTTTTTAAA GTATAAATAA ATCTCTATTT CTTGCGACCT  
 3910 3920 3930 3940 3950 3960  
 ACATTGAACT TAATAGTTAT GTTTTACAGA GGATCACTTC ATCGAAACGG GCCACCATCT  
 3970 3980 3990 4000 4010 4020  
 CCATTAAACC TTCAAATGAC TACATTTGAT TCGGAGATCC CTCTGACTTT TGATCAGGTG  
 4030 4040 4050 4060 4070 4080  
 AGTCTTACAT TGAGTTCAAA CTTTTTGAAT TTAAGCGTTC TATCTGATAA AGTTCTTCGG  
 4090 4100 4110 4120 4130 4140  
 TGGTTTTATA ATTTTGTATT CATAAACTTA CCCACTCCTT TCTCACTAAC ATTTTACCCT  
 4150 4160 4170 4180 4190 4200  
 GTTCAGCTGC CAGTTTCCAT GGAATCCGAT AGACCCCTGA TTGAAGAGGT AACTGTGAAA  
 4210 4220 4230 4240 4250 4260  
 GTAGTCAATT AATTCCCTGT GTTTCTACCC CACTCAATCC TTTTGTATTT TTTGTTTCAGT  
 4270 4280 4290 4300 4310 4320  
 CTATCCACTA TCAATGTCTT ATCACCTCTA GATACTGTTT AGAAGAAAAT ATTGTTTACA  
 4330 4340 4350 4360 4370 4380  
 GTTATGGAAA TCACATATAC TTTGTTCTGG AATTGTATAT GTATGCTTTG AAAAAGCACA  
 4390 4400 4410 4420 4430 4440  
 TTAGAATACT ACAAACATTA GTTTCCATCA GATTTTTGAT TTATCAAAAC CGTTATATTA  
 4450 4460 4470 4480 4490 4500  
 GACACTCTTA AGTTATCATA TTCTAATTC CAAGAATGTT ATATTTTGAA GAAGCCGGTG  
 4510 4520 4530 4540 4550 4560  
 ATTGTCAAAA AGATTGAAAA CTCCGAGTTT CTATATATGC GAAATTTTCA CTTCAGCCCA

4570            4580            4590            4600            4610            4620  
 CACACACACA CACACATTCA CGAAACTTTG TGTGTTTAT GTTACTTATA TGTATCTTT

4630            4640            4650            4660            4670            4680  
 TCTGTCTGAT CATGGTTTTT GGAAGTATA TGTGAGCCAC

4690            4700            4710            4720            4730            4740  
 ATTGATTAAA CCTGTGAGAG ATGCCCATTT GTACTCATTT TACGACTGTC TCATGTCCAA

4750            4760            4770            4780            4790            4800  
 ACACCATGTT TATTGTAATT ACCAGGCTAC TATTTGCAGA TGCATCAAC ATCACCACCT

4810            4820            4830            4840            4850            4860  
 CCACCATCTG GATGTCTGGC CAGATTCCAT CCGAAGCAG TGGACAAATT CTCCATTGTA

4870            4880            4890            4900            4910            4920  
 GCTTTTCCAT TGGCATTTAC AATGTTTAAAT GTTAGTTAAT CCACAGTTAA AAATTCCCAT

4930            4940            4950            4960            4970            4980  
 AATCATAAAT ATCTCGACTT TTCAGCTTGT CTAAGTGGT CACTATTTGT CTCAACTTT

4990            5000            5010            5020            5030            5040  
 CGATCAAAAC TATCAGTGAT TGAAGTTTAT CCCTTTTAAAT TCCAATAATT CACAGTTGCC

5050            5060            5070            5080            5090            5100  
 GGTATCTACC TCCATTCTTT TCCGATGATT CGCAGTTTTT CACAGGGTTC AAATGTATCT

5110            5120            5130            5140            5150            5160  
 CGTTCAATCT TTTTATGGTT ATTTCTCTTG AATGTCCATT TTAATATTTA TAGAACAATT

5170            5180            5190            5200            5210            5220  
 TTATGTACAT TGTGTTGGTA TTCAATTCGA AAAACAATGA AATTTATTTT TAAATAACTG

5230            5240            5250            5260            5270            5280  
 CGTTTCTGGG GTTTCTATCA GCACTTACTA GCTGACAAAA ACTTTTCCGT ATTCCGAATT

5290            5300            5310            5320            5330            5340  
 AGATTTTAT GCAAGCAATG TTTCATTTT ACACAGTATA GTATTTATTC TTACTTTTGA

5350            5360            5370            5380            5390            5400  
 TTATATTGCT CGCACCTAA ATGACAGGTA TTAGAAATTA ACCGCTTTTC AGAGTATTTT

5410            5420            5430            5440            5450            5460  
 TAATCTTCTT AGTACTAGTT TAGTTCTTTA AATAAGAAAC CATCTAGTTT TTCATTATCA

5470            5480            5490            5500            5510            5520  
 CTCAACTTCA GTCGGACAAA TTTTAAATTT TTTACTCGAT AAAAAAATTT TATAATTTCAG

5530            5540            5550  
 ACAAATTATG TCTTCTCATT TTTGATCGCT



10 20 30 40 50 60  
 ATGAAGTTTA TTCCTGAAAT CACACTACTC TTGCTTTTAT TTGTACACTC TACACAGGCT  
 70 80 90 100 110 120  
 AAAGGAAAAC GACGGAAATG TCCGGAGGGT GCGTGGTCGG AAGGAAAGAT TATGAACACG  
 130 140 150 160 170 180  
 ATCATGAGCA ACTACACGAA AATGTTGCCC GACGCGGAGG ACAGCGTACA AGTTAATATT  
 190 200 210 220 230 240  
 GAGATTCATG TACAGGATAT GGAAGCTTG AATGAAATAT CATCCGACTT TGAAATTGAC  
 250 260 270 280 290 300  
 ATTTTATTCA CTCAACTGTG GCATGACTCG GCACTTTCTT TTGCTCATCT TCCGGCTTGT  
 310 320 330 340 350 360  
 AAGCGAAATA TCACAATGGA AACACGACTT TTACCTAAGA TTTGGTCTCC AAACACGTGT  
 370 380 390 400 410 420  
 ATGATTAATT CAAAACGAAC AACCGTCCAT GCATCACCAT CGGAAAATGT GATGGTTATT  
 430 440 450 460 470 480  
 CTGTACGAGA ATGGAACAGT CTGGATTAAC CATCGTCTTA GTGTCAAATC ACCTTGCAAT  
 490 500 510 520 530 540  
 TTGGATCTGC GACAGTTTCC TTTCGATACT CAAACTTGCA TATTAATCTT TGAATCCTAT  
 550 560 570 580 590 600  
 AGTCATAACT CAGAAGAAGT TGAACCTCAT TGGATGGAAG AAGCTGTCAC ATTAATGAAG  
 610 620 630 640 650 660  
 CCAATTCAAC TTCCTGACTT TGATATGGTT CATTATTCAA CTAAAAAGGA AACTTTACTC  
 670 680 690 700 710 720  
 TATCCAAACG GGTACTGGGA TCAGCTTCAA GTTACTTTCA CTTTCAAACG ACGATATGGA  
 730 740 750 760 770 780  
 TTCTATATTA TTCAAGCCTA TGTTCACAAC TATCTTACAA TCATTGTATC TTGGGTTTCA  
 790 800 810 820 830 840  
 TTCTGCATGG AACCAAAAGC TCTGCCGGCA AGAACAACCTG TCGGAATCTC ATCTCTTCTA  
 850 860 870 880 890 900  
 GTTCTTACTT TCCAGTTTGG AAATATTTTG AAAAATCTTC CAAGGGTTTC ATATGTGAAA  
 910 920 930 940 950 960  
 GCAATGGATG TGTGGATGCT TGGATGCATA TCATTTGTCT TCGGAACCAT GGTAGAATTG  
 970 980 990 1000 1010 1020  
 GCATTTGTTT GTTACATTTT CCGTTGTCAG AACAGCGTAA GAAACGCGGA ACGACGACGG  
 1030 1040 1050 1060 1070 1080  
 GAACGAATGA GAAATTCTCA GGTGTGGGCA AACGGATCGT GTAGAACTAG AAGCAACGGG  
 1090 1100 1110 1120 1130 1140  
 TATGCAAACG GGGGATCTGT AATCTCACAT TATCATCCAA CAAGCAATGG AAATGGGAAT

Fig. 8  
Page 1

1150	1160	1170	1180	1190	1200
AATAATCGAC	ATGATACACC	TCAAGTTACT	GGAAGAGGAT	CACTTCATCG	AAACGGGGCCA
1210	1220	1230	1240	1250	1260
CCATCTCCAT	TAAACCTTCA	AATGACTACA	TTTGATTCCG	AGATCCCTCT	GACTTTTGAT
1270	1280	1290	1300	1310	1320
CAGCTGCCAG	TTTCCATGGA	ATCCGATAGA	CCCCTGATTG	AAGAGATGCG	ATCAACATCA
1330	1340	1350	1360	1370	1380
CCACCTCCAC	CATCTGGATG	TCTGGCCAGA	TTCCATCCGG	AAGCAGTGGA	CAAATTCTCC
1390	1400	1410	1420	1430	1440
ATTGTAGCTT	TTCCATTGGC	ATTTACAATG	TTTAATCTTG	TCTACTGGTG	GCACTATTTG
1450	1460	1470			
TCTCAAACCTT	TCGATCAAAA	CTATCAGTGA			

002240" 2296560






Fig. 9

09599622.043700

# MOD-1 Selectivity




Fig. 10




Figure 11

090909 042700

04/01/2000

COMBINED DECLARATION AND POWER OF ATTORNEY

As a below named inventor, I hereby declare that:

My residence, post office address and citizenship are as stated below next to my name.

I believe I am the original, first and sole inventor (if only one name is listed below) or an original, first and joint inventor (if plural names are listed below) of the subject matter which is claimed and for which a patent is sought on the invention entitled A NOVEL SEROTONIN-GATED ANION CHANNEL, the specification of which

☒ is attached hereto.

☐ was filed on \_\_\_\_\_ as Application Serial No. \_\_\_\_\_  
and was amended on \_\_\_\_\_.

☐ was described and claimed in PCT International Application No. \_\_\_\_\_  
filed on \_\_\_\_\_ and as amended under PCT Article 19 on \_\_\_\_\_.

I hereby state that I have reviewed and understand the contents of the above-identified specification, including the claims, as amended by any amendment referred to above.

I acknowledge the duty to disclose all information I know to be material to patentability in accordance with Title 37, Code of Federal Regulations, §1.56(a).

**FOREIGN PRIORITY RIGHTS:** I hereby claim foreign priority benefits under Title 35, United States Code, §119 of any foreign application(s) for patent or inventor's certificate or of any PCT international application(s) designating at least one country other than the United States of America listed below and have also identified below any foreign application for patent or inventor's certificate or any PCT international application(s) designating at least one country other than the United States of America filed by me on the same subject matter having a filing date before that of the application(s) of which priority is claimed:

Country	Serial Number	Filing Date	Priority Claimed?
			Yes/No

**PROVISIONAL PRIORITY RIGHTS:** I hereby claim priority benefits under Title 35, United States Code, §119(e) and §120 of any United States provisional patent application(s) listed below filed by an inventor or inventors on the same subject matter as the present application and having a filing date before that of the application(s) of which priority is claimed:

Serial Number	Filing Date	Status
60/131,149	April 27, 1999	Pending

## COMBINED DECLARATION AND POWER OF ATTORNEY

**NON-PROVISIONAL PRIORITY RIGHTS:** I hereby claim the benefit under Title 35, United States Code, §120 of any United States application(s) listed below and, insofar as the subject matter of each of the claims of this application is not disclosed in the prior United States application in the manner provided by the first paragraph of Title 35, United States Code, §112, I acknowledge the duty to disclose all information I know to be material to patentability as defined in Title 37, Code of Federal Regulations, §1.56(a) which became available between the filing date of the prior application and the national or PCT international filing date of this application:

Serial Number	Filing Date	Status

I hereby appoint the following attorneys and/or agents to prosecute this application and to transact all business in the Patent and Trademark Office connected therewith: Paul T. Clark, Reg. No. 30,162, Karen L. Elbing, Ph.D., Reg. No. 35,238, Kristina Bieker-Brady, Ph.D., Reg. No. 39,109, Susan M. Michaud, Ph.D., Reg. No. 42,885, Mary Rose Scozzafava, Ph.D., Reg. No.36,268, James D. DeCamp, Ph.D., Reg. No. 43,580.

Address all telephone calls to: Kristina Bieker-Brady, Ph.D. at 617/428-0200.

Address all correspondence to: Kristina Bieker-Brady, Ph.D. at Clark & Elbing LLP, 176 Federal Street, Boston, MA 02110.

I hereby declare that all statements made herein of my own knowledge are true and that all statements made on information and belief are believed to be true; and further that these statements were made with the knowledge that willful false statements and the like so made are punishable by fine or imprisonment, or both, under Section 1001 of Title 18 of the United States Code and that such willful false statements may jeopardize the validity of the application or any patents issued thereon.

Full Name (First, Middle, Last)	Residence Address (City, State, Country)	Post Office Address (Street, City, State, Country)	Citizenship
Rajesh Ranganathan	Cambridge, MA	387 Columbia Street Cambridge, MA 02141	India
Signature:			Date:

### COMBINED DECLARATION AND POWER OF ATTORNEY

Full Name (First, Middle, Last)	Residence Address (City, State, Country)	Post Office Address (Street, City, State, Country)	Citizenship
H.R. Horvitz	Auburndale, MA	54 Maple Street Auburndale, MA 02166	United States
Signature:			Date:

Full Name (First, Middle, Last)	Residence Address (City, State, Country)	Post Office Address (Street, City, State, Country)	Citizenship
Stephen C. Cannon	Needham, MA	88 Greendale Avenue Needham, MA 02494	United States
Signature:			Date: